



**EFFECTS OF *EUPATORIUM ODORATUM* LEAF MEAL ON
GROWTH PERFORMANCE, HAEMATOLOGY, SERUM
BIOCHEMISTRY AND CARCASS CHARACTERISTICS OF
BROILER FINISHER BIRDS**

Igugo, R. U¹., Madu, P. O²., Odo, B.I¹,

¹Department of Animal/Fisheries Science and Management, Enugu State
University of Science And Technology, ESUT.

²Animal Science Dept, University of Nigeria Nsukka

*Author for Correspondence: igugo@esut.edu.ng

ABSTRACT

The study was conducted to investigate the effects of *Eupatorium odoratum* leaf meal (EOLM) on growth performance, carcass characteristics, haematology and serum biochemical indices of broiler finisher birds. A total of one hundred and forty-four (144) day-old Ross strain commercial broiler chicks were used for the study. The birds were randomly allotted to four dietary treatments (T₁, T₂, T₃ and T₄) with 36 birds per treatment. Each treatment was further replicated three times with 12 birds per replicate in a Completely Randomized Design (CRD). T₁, T₂, T₃ and T₄ represent 0mg, 100mg, 200mg and 300mg inclusion levels of EOLM/kg of diet, respectively. The experimental diets and water were provided *ad libitum* throughout the 8 weeks' period of the study. Data were collected on growth performance (feed intake, live weight, live weight gain, FCR and percentage mortality), carcass characteristics, haematology and serum biochemical indices. All data were subjected to Analysis of Variance using statistical product and service solution (Version 23) while significant treatment means were separated using Duncan's Multiple Range Test at 5% level of significance. Results of growth performance showed that there were significant differences (p<0.05) in the average daily weight gain, feed conversion ratio and mortality percentage. T₃ and T₄ had the highest average weight gain (86.04g±1.11 and 86.15g±1.22) followed by T₁ (85.92g±1.11) with lowest from T₂ (79.61g±1.01). There were no significant differences (p>0.05) in feed intake among the treatment groups. FCR was best in T₃ (1.82±0.12) followed by T₁ (1.88±0.01) and worst in T₂ (2.07±0.04). Mortality percentage was highest in T₁ (6.03%±0.03) but lowest in T₃/T₄ (3.4%±0.05/2.68%±0.13). Results of haematological indices were also significantly (p<0.05) affected by EOLM. PCV values were statistically similar in T₂ (31.90%±0.04) and T₃ (31.70%±1.33) both of which were higher than T₄ (28.80%±2.19). PCV value in T₄ did not vary with T₁ (25.65%±0.49). Haemoglobin values were similar in T₁ (9.27±0.32), T₂ (10.88±0.11) and T₄ (10.05±0.59). These values however, were significantly lower (p<0.05) than T₃ (11.63±0.73). TWBC was significantly higher in T₁ (52.50±6.64), T₂ (42.23±1.01) and T₄ (43.00±1.73) than in T₃ (80.75±0.43). Monocyte values were higher in T₁ (4.50±0.29), T₃ (5.00±1.73) and T₄ (8.00±1.15) than T₂ (3.50±0.87). However, RBC, lymphocyte, and heterophils (%) values did not significantly differ (p>0.05) among the treatment groups. Results on serum biochemistry showed that albumin and alkaline phosphatase were significantly (p<0.05) affected by EOLM supplementation. Albumin value of T₂ (28.18±0.09) was higher than values from the other groups (T₁ [26.75±0.04], T₃ [27.33±0.29] and T₄ [26.82±0.41]). Alkaline phosphatase level (ALP) was highest in T₃/T₄ (122.64±3.45/127.45±2.79) followed by T₁ (102.42±5.10) and the least in T₂ (82.85±9.15). Results on carcass characteristics showed that T₃ (2577.24±1.50) had the highest live weight (g) followed by T₄/T₁ (2571.34±0.57/2566.01±1.21) and the least value in T₂ (2384.44±0.57). De-feathered weight, dressed weight and dressing percentage had similar trend of values. However, there were no significant differences (p>0.05) in breast weight (%), back cut (%), thigh (%), drumstick (%), kidney (%), heart (%), spleen (%) and liver (%). It can be concluded that dietary inclusion of EOLM at 200 mg/kg diet improved growth performance, haemato-biochemical indices in broiler finisher chickens.

INTRODUCTION

Nigeria is one of the lowest animal protein consumption nation in the world, with the diet of an average Nigerian containing lower than recommended requirement by the Food and Agricultural Organisation (Egugozie, 2019). In order to accelerate nutritional well-being and economic development in Nigeria, the supply of meat must be significantly boosted through livestock farming in the next years. Food insecurity is quickly rising in some third world countries and achieving relatively self-sufficient animal protein supply

necessitates intensive production of meat animals, particularly poultry, which is now produced at a small-scale subsistence levels (Wong et al. 2017). However, Alemu (2020) reported that malnutrition is significantly connected to poor meat consumption, particularly among children in developing nations. Several studies recommend a supply-side approach to meet consumption needs, such as increasing herd size, livestock density and supply chain efficiency, while noting the negative impacts on land and environment as well (Godde et al. 2021).

The Nigerian poultry industry comprises

approximately 180 million birds and produces 300 million tons of meat and 650 million tons of eggs per year. Yet, local production only meets 30% of the demand for chicken eggs and meat. Thus, there is plenty of room for growth in the industry most of which are domiciled in the southern part of the country either in semi-intensive farms or intensive ones (FAO, 2018; Mba, 2021). Egg and meat production are major divisions of poultry production (United States Department of Agriculture – USDA, 2018). Broiler birds are those kept and reared for meat production from day-old to about eight weeks of age for good quality tender meat as source of protein in human diet. However, the profit levels in this farming enterprises have been constrained by a number of factors critical among which is feed quality, mortality and general cost incurred. The high feed cost is partly due to competition for limited grains between humans and animals. Feed cost as a main component of input costs and profitability in broiler production necessitates a thorough examination. Farmers have been progressively designing poultry meals using locally accessible feed ingredients in recent years (Escobedo del Bosque et al., 2020). Previously, broiler meat is relatively inexpensive, high in nutrients, culturally and religiously acceptable, and has a high output turnover rate (Nemeth et al. 2019). But in recent years, in some developing countries like Nigeria, it is becoming expensive.

Poultry production is undergoing a continuous challenge to develop management strategies to optimize chickens' efficiency while limiting food safety concerns (Abdelli et al. 2021). Traditionally, antimicrobials have been widely used for improving health and growth performance in poultry; however, the increased public awareness about the risk of developing cross resistance of pathogens to antibiotics has resulted in the gradual removal of antibiotics for therapeutic and prophylactic uses in food animals (Ricke et al. 2020). The shift away from antibiotic supplementation has resulted in a tremendous growth in research focusing on the implementation of effective alternative control methods, management and dietary amendments aiming to improve animal health, welfare, and productivity (Abdelli et al. 2021). A wide range of feed additives including a broad spectrum of essential oils and related compounds from botanical sources to organic acids (Zhai et al. 2018), as well as probiotics and prebiotics (Al-Khalafah, 2018), chemicals such as aldehydes (Ricke et al. 2019), bacteriophages (Li et al., 2020), zinc oxide (Swain et al. 2016), exogenous enzymes (Torres-Pitarch et al. 2019) and competitive exclusion products (Schneitz and Hakkinen, 2016) have been used in animal production.

Phytogenics such as *Eupatorium odoratum* with several biological properties have favourable influence on the health and growth of poultry (Gopi et al. 2018), and are considered a modern class of feed additives that have received widespread interest in the chickens

industry (Abd El –Hack et al. 2016). Plants and herbal extracts contain many bioactive substances like alkaloids, flavonoids, soaps, phenolic substances and polypeptides (Al-Yasiry and Kiczorowska, 2016). The mechanism of action of these compounds was different due to the multiplicity of their types as they have antibacterial effects, immune stimuli, antioxidants and growth stimuli (Abd El-Hack and Alagawany, 2015).

In evaluating these feed resources, it is important to check its effect on the health status of the livestock (Vastolo et al. 2022). In recent times, there is global utilization of herbal medicine in the treatment of various diseases affecting both humans and animals (Megersa and Woldetsadik, 2022). *Eupatorium odoratum* is one of these medicinal plants with overemphasized values. This plant may play a role in improving production and immune system of birds against different diseases (Jha et al. 2022). Hence, with strong medicinal values, it could be effectively utilized as natural growth promoter, replacing antibiotics and other synthetic feed additives (Tiamiyu et al. 2021). Anti-inflammatory properties of *E. odoratum* is known among herbaceous plants because it possesses chemical compounds with membrane stabilizing property (Toghyani et al. 2011). Nevertheless, it is expected that compounds with membrane stabilizing properties should proffer protection to cell membranes against injurious substances and thereby exhibit anti-inflammatory properties (Akhter et al. 2022).

Location and Duration of the Experiment

This experiment was conducted at the Poultry Unit of the Department of Animal Science Teaching and Research Farm, University of Nigeria, Nsukka. Nsukka is situated at the intersection of Longitude 7° 12.5' N and Latitude 6° 45' E in the derived Savannah area (Breinholt et al. 1981), with a maximum plateau altitude of 443m and a minimum of 207m above sea level (Ofamata, 1978). However, Breinholt et al. (1981) reported that the area has a relative humidity range of 73.1 percent to 76.6 percent, the climate is humid tropical and range of annual rainfall of 1680mm to 1700mm. The study lasted through a period of 8 weeks (56 days).

Management of Experimental Birds

A total of one hundred and forty-four (144) day old broiler birds were used for the experiment. The research contained 4 different treatments replicated thrice and each replicate contained 12 birds; making it 36 birds per treatment. Before the arrival of the birds, the pens measuring 12x10m² divided into different cells were cleaned and disinfected. The birds were brooded for 4 weeks before application of the test ingredient (finisher phase) so as to get the birds acclimatized to the environment and also escape brooding stress. Their

initial body weight was recorded before being randomly placed in deep litter pens. *Ad-libitum* feed and water (containing routine vaccines/medications) were provided to the birds under the same management conditions.

The Experimental Diet and Treatments

Fresh leaves of *Eupatorium odoratum* were harvested within the campus of University of Nigeria, Nsukka, Enugu State, Nigeria, defoliated from the stem and air-dried under room temperature. The air-dried leaves were ground into powder and sieved to become meal prior to storage for subsequent use to prepare broiler

feed at various levels of inclusion. The product was incorporated into a commercial feed sourced from an Olam Company (Chikun Feed: Broiler Finisher diet) with the proximate composition outlined in Table 7.

The following were the treatment groups:

- Ø Treatment 1: 0mg of *E. odoratum* leaf meal per kg diet.
- Ø Treatment 2: 100mg of *E. odoratum* leaf meal per kg diet.
- Ø Treatment 3: 200mg of *E. odoratum* leaf meal per kg diet.
- Ø Treatment 4: 300mg of *E. odoratum* leaf meal per kg diet.

Table 1: Proximate Composition of the Experimental rations.

Composition	Broiler Starter Diet	Broiler Finisher Diet
Crude fiber (max)	5.0 %	5.0 %
Crude protein (min)	21.0 %	18.0 %
Crude fat (min)	4.0 %	5.0 %
Calcium (min)	1.0 %	0.9 %
Available phosphorus (min)	0.5 %	0.4 %
Lysine (min)	1.2 %	0.9 %
Methionine (min)	0.5 %	0.4 %
Metabolizable Energy (min)	2900 kcal/kg	3000 kcal/kg

*As labelled by the manufacturers.

Data Collection Data collection were carried out based on the specific objectives of the study for growth performance, haematological indices, serum biochemistry, carcass and organ characteristics.

Growth Performance: Birds were weighed at day old using a sensitive scale and subsequently on a weekly basis till 8weeks of age. From the weekly body weight, body weight gain was calculated as the final weight minus the initial body weight.

Average Daily Weight Gain (kg): This was the weight gained by the birds in each treatment at the end of the study and it was obtained as:

$$\frac{\text{Final weight -Initials weighs of birds /treatment}}{\text{Number of birds}}$$

Average Daily Feed Intake (g): This is the total amount of feed in g consumed per bird per treatment obtained as:

$$\frac{\text{Feed supplied - Feed leftover/treatment}}{\text{Number of birds}}$$

Feed Conversion Ratio: This is a measure of the efficiency at which the broilers in each treatment converted feed into desired output (flesh) and it is equal to

$$\frac{\text{Feed Intake/treatment}}{\text{Weight Gain}}$$

Mortality (%): This is the number of animals that died in each treatment during the experiment and expressed in percent calculated as:

$$\frac{\text{Number of dead birds/treatment}}{\text{Number of Live Birds}} \times 100$$

Haematological Indices: At the end of 8 weeks of the experiment, 6 birds were selected randomly per treatment group and subjected to blood sample collection for the determination of haematological indices. Data were collected for; red blood cell count, total white blood counts, packed cell volume, haemoglobin (Hb), lymphocytes, heterophils, eosinophils and monocytes. Blood samples were collected from jugular vein of sampled birds into 1ml vials containing 2mg ethylene-diamine tetra-acetic acid (EDTA). The tests were done with the help of an automatic haematology analyzer as reported by Aikpitanyi and Egweh (2020). Data were recorded in their relevant units of measurement.

Serum Biochemistry: Jugular vein piercing was used to obtain blood samples from 6 birds taken randomly from each group at the end of the experiment and placed in containers without anticoagulant. Following collection, the samples were centrifuged for 15 minutes at 1500 rpm to sediment the blood cells and the supernatant frozen for serum biochemical analysis. Parameters considered were: total protein, albumin, uric acid, globulin, aspartate aminotransferase (AST), alanine aminotransferase (ALT) and alkaline phosphatase (ALP). These were carried out according to the methods described by Bahman *et al.* (2011).

Carcass and Organ Characteristics: At the end of the experiments, one bird per replicate was randomly selected, starved of feed for 12 hours, and slaughtered for carcass evaluation. Each bird was scalded and defeathered for carcass and organ weight determination. Each part of chicken was weighed on a sensitive scale and the weight recorded in gram. Dressing percentage was calculated as the percentage of the carcass weight to the final live weight. The internal organ weights and weights of

the cut parts were expressed as percentage of carcass weight. These procedures were carried out according to the methods reported by Jiwuba *et al.* (2018) and Ukam *et al.* (2020). Parameters considered were as follows; live weight, dressed weight, de-feathered weight and percentage dressing, breast cut, back cut, thigh, drumstick, kidney, heart, spleen and liver. All carcass measurement values were recorded in gram and percentages, respectively.

Experimental Design

The experiment were conducted in a completely randomized experimental design (CRD) and the mathematical model used is presented below:

$$Y_{ij} = \mu + X_i + \Sigma_{ij}$$

Where:

Y_{ij}	=	Individual observations
μ	=	Population means
X_i	=	Treatment effect (effects of <i>E. odoratum</i>)
Σ_{ij}	=	Experimental error

3.6 Statistical Analysis

The data were subjected to analysis of variance (ANOVA) and significantly different means were separated using Duncan's New Multiple Range Test on SPSS version 23 (2015).

RESULTS AND DISCUSSION

Growth Performance of Broiler Finisher Birds fed Diets Containing *Eupatorium odoratum* Leaf Meal

Results of the growth performance of broiler birds fed diets supplemented with *Eupatorium odoratum* are shown in Table 8.

Table 2: Growth performance of broiler birds fed diets containing *Eupatorium odoratum* leaf meal.

Parameters	Treatment 1 (control)	Treatment 2 (100mg <i>E.O</i>)	Treatment 3 (200mg <i>E.O</i>)	Treatment 4 (300mg <i>E.O</i>)	p. value
IBW (g)	160.31±1.21	155.49±1.21	168.01±1.24	159.11±1.07	0.21
FBW (g)	2566.01±1.21 ^b	2384.44±1.04 ^c	2577.24±1.50 ^a	2571.34±0.57 ^b	0.00**
TWG (g)	2405.70±1.01 ^b	2228.95±1.51 ^c	2409.23±1.09 ^a	2412.23±1.70 ^a	0.00**
ADWG (g)	85.92±1.11 ^b	79.61±1.01 ^c	86.04±1.11 ^a	86.15±1.22 ^a	0.01**
TFI (g)	4531.01±0.11	4610.12±0.41	4392.51±0.40	4741.22±0.44	0.71 ^{NS}
ADFI (g)	161.82±0.06	164.65±0.06	156.45±0.07	169.33±0.09	1.00 ^{NS}
FCR	1.88±0.11 ^c	2.07±0.04 ^a	1.82±0.12 ^d	1.97±0.11 ^b	0.01**
Mortality (%)	6.03±0.03 ^a	5.15±0.71 ^a	3.41±0.05 ^b	2.68±0.13 ^b	0.00**

^{abcd}: Means on the same row with different superscripts are significantly different ($P < 0.05$), IBW; Initial body weight, FBW; Final body weight, TWG; Total weight gain, ADWG; Average daily weight gain, TFI; Total feed intake, ADFI; Average daily feed intake, FCR; feed conversion ratio.

Results of the growth performance shows that there were significant differences ($P < 0.05$) in the final body weight (FBW), average daily weight gain (ADWG), feed conversion ratio (FCR) and mortality percentage. Treatment 3 recorded highest final body weight value of 2577.24±1.50g whereas the lowest value was recorded in treatment 2 (2384.44±1.04g). Treatment 3 and 4 had the highest average weight gain (86.04±1.11g and 86.15±1.22g, respectively) followed by treatment 1 (85.92±1.11g) with the lowest value from treatment 2 (79.61±1.01g). FCR was highest in treatment 3 (1.82±0.12) followed by treatment 1 (1.88±0.11) and lowest in treatment 2 (2.07±0.04). Mortality percentage was highest in treatment 1 (6.03±0.03%) but lowest in treatment 3 and treatment 4 (3.41±0.05% and 2.68±0.13%, respectively). Feed intake was however not affected ($P > 0.05$) by the test ingredient.

The significant differences obtained in FBW and FCR were similar to the reported values of Jiwuba et al. (2018) and Ekwe et al. (2017) but values for weight gain and mortality percentage differed from those of the authors. Thus, those observed differences in this study could be as a result of source of the *E. odoratum*, methods of processing or prevailing environmental conditions. However, the results of this study are in contrast with the report of Donkoh et al. (2002) which indicated that, the inclusion of *E. odoratum* leaf meal had an adverse effect on the performance of broiler chickens by reducing feed intake, body weight gain, feed conversion efficiency which could be as a result of higher percentage of tannin present in *E. odoratum*. Improved body weight gain recorded in this study

could be attributed to the fact that *E. odoratum* leaf has a good potential for feeding of livestock due to its high crude protein level, and low extractible phenolic content (Apori et al. 2000). Thus, this is in agreement with the results of Bamikole et al. (2004) who reported good performance for the weight gain at 30% dietary level of *E. odoratum* leaf meal. This result is also in line with the report of Alabi and Chime (2008) which showed that inclusion of *E. odoratum* leaf meal in layers mash significantly ($P < 0.05$) improved FCR. The presence of water infusion of *E. odoratum* leaf meal may have initiated enzymatic activities in the gastrointestinal tract thereby improving digestion and assimilation of nutrients (Ekwe et al. 2017). It was observed that reduction in mortality occurred when birds were fed *Eupatorium odoratum* at levels of 200 to 300mg and this corroborates the results of the study by Amaliah et al. (2019). The research work of Alisi et al. (2011) and Marianne et al. (2014), explained that *E. odoratum* contains antioxidants and is able to prevent free radicals which are believed to be the cause of various degenerative disease and organ damages that may be responsible for increased mortality in the chicken.

Results of the Haematological Indices of Broiler Finisher Birds fed Diets Containing *Eupatorium Odoratum* Leaf Meal.

The results of the haematological indices of broiler birds fed diets containing *Eupatorium odoratum* leaf meal are presented in Table 9.

Table 3: Haematological indices of broiler birds fed *Eupatorium odoratum* leaf meal

Parameters	Treatment 1 (control)	Treatment 2 (100mg <i>E.O</i>)	Treatment 3 (200mg <i>E.O</i>)	Treatment 4 (300mg <i>E.O</i>)	p. value
Packed Cell Volume (%)	25.65±0.49 ^b	31.90±0.40 ^a	31.70±1.33 ^a	28.80±2.19 ^{ab}	0.03*
Haemoglobin (g/dl)	9.27±0.32 ^b	10.88±0.11 ^{ab}	11.63±0.73 ^a	10.05±0.59 ^{ab}	0.04*
RBC (x10 ⁶ /μl)	2.41±0.01	2.95±0.05	3.19±0.20	2.96±0.32	0.09 ^{NS}
TWBC (x10 ³ /μl)	52.50±6.64 ^b	42.25±1.01 ^b	80.75±0.43 ^a	43.00±1.73 ^b	0.00**
Lymphocytes (%)	55.50±7.79	46.00±1.15	51.50±0.29	61.00±4.04	0.18 ^{NS}
Heterophils (%)	30.50±6.06	26.50±0.29	33.50±3.18	22.50±6.06	0.40 ^{NS}
Eosinophils (%)	7.50±0.87	9.50±0.29	10.00±1.15	8.50±0.87	0.24 ^{NS}
Monocytes (%)	4.50±0.29 ^b	13.50±0.87 ^a	5.00±1.73 ^b	8.00±1.15 ^b	0.00**

^{ab}: Means on the same row with different superscripts are significantly different (P<0.05), RBC: red blood cell count; TWBC: total white blood cell count.

Results of some haematological indices were also significantly (P<0.05) affected by *E. odoratum* leaf meal. For instance, PCV values were statistically similar in treatment 2 (31.90±0.40%) and treatment 3 (31.70±1.33%), both of which were significantly (P<0.05) higher than treatment 4 (28.80±2.19%). However, the value in treatment 4 did not differ from that of treatment 1 (25.65±0.49%). Hb values were higher in treatment 2 (10.88±0.11 g/dl), treatment 3 (11.63±0.73 g/dl) and treatment 4 (10.05±0.59 g/dl) compared to treatment 1 (9.27±0.32 g/dl). TWBC count was significantly higher in treatment 1 (52.50±6.64 x 10³/μl), treatment 2 (42.25±1.01 x 10³/μl) and treatment 4 (43.00±1.73 x 10³/μl) and highest in treatment 3 (80.75±0.43 x 10³/μl). Monocyte values were higher in treatment 1 (4.50±0.29%), treatment 3 (5.00±1.73%) and highest in treatment 4 (8.00±1.15%). However, lymphocyte, heterophils and eosinophil (%) values did not vary among the treatment groups.

Haematological indices are often used for the diagnosis and treatment of animal diseases. The values (Table 9) obtained in this study were within normal ranges for broilers signifying that there were no symptoms of physiological anemia in the experimental birds (Al-Nedawi, 2018). The haemoglobin and PCV values were higher (P<0.05) in broilers supplemented with 200g *Eupatorium odoratum*. Haemolysis results from acute inflammation from most pathogenic microorganisms which is manifested in lower haemoglobin level and PCV (Osho et al., 2014). Thus, the improved values in birds fed with 200mg *Eupatorium odoratum* could be due to the inability

of the organisms to cause haemolysis, resulting from the anti-inflammatory potentials inherent thereof. This improvement could also be associated with a better utilization of feed which facilitates high nutrients absorption that leads to higher haemoglobin concentration and higher percentage of PCV (Ekwe et al., 2017). More so, the results of this study suggest the potential of *Eupatorium odoratum* to act as an immuno-stimulant through the alteration of the occurrences and distribution of the white blood cell counts and the differentials. The neutrophils and lymphocytes percentage could be an indication of improved immune response (Osho et al. 2014). Dhabhar (2000) had stressed that higher lymphocyte and WBC counts are associated with the ability of animals to perform well under a very stressful condition. Furthermore, it means that the implication of the TWBC of birds recording 80.75 x 10³/μl in treatment 3 showed a high immunity against infections. The non-significant effect of the leaf meal on the eosinophil count of the birds was an indicator that the anti-nutritional factors present in *Chromolaena odorata* leaf did not affect the blood quality of the birds (Okosun and Oyedeji, 2016). Other reports by Okwesili et al. (2021), Machebe et al. (2011) and Olabatoke and Oloniruha (2009) shows that antioxidant phytochemicals play a protective role on the lymphocytes and reduce their destruction in the blood.

Serum Biochemistry of Broiler Finisher Birds fed Diets Containing *Eupatorium Odoratum* Leaf Meal

Results of the serum biochemistry of broiler birds on diets supplemented with *Eupatorium odoratum* leaf meal are shown in Table 10.

Table 4: Serum Biochemistry of broiler birds fed diets containing *Eupatorium odoratum* leaf meal.

Parameters	Treatment 1 (control)	Treatment 2 (100mg <i>E.O</i>)	Treatment 3 (200mg <i>E.O</i>)	Treatment 4 (300mg <i>E.O</i>)	p. value
TP (g/l)	37.11±2.22	38.74±2.17	39.85±2.81	32.10±1.38	0.14 ^{NS}
ALB (g/l)	26.75±0.04 ^b	28.18±0.09 ^a	27.33±0.29 ^b	26.82±0.41 ^b	0.01 ^{**}
Uric acid (µmol/l)	231.03±41.16	318.40±23.52	196.19±29.71	187.07±37.45	0.08
ALP (Iµ/L)	102.40±5.10 ^c	82.85±9.15 ^b	122.64±3.45 ^a	127.45±2.79 ^a	0.00 ^{**}
ALT (Iµ/L)	10.09±0.28	9.61±0.15	9.54±0.05	9.87±0.11	0.16 ^{NS}
AST (Iµ/L)	64.90±13.11	83.74±3.82	88.84±0.56	74.01±11.04	0.29 ^{NS}
Globulin (g/l)	10.37±2.18	10.56±2.25	12.53±3.10	5.28±1.79	0.24 ^{NS}

^{abc}: Means on the same row with different superscripts are significantly different (P<0.05), ALT: alanine transaminase; AST: aspartate transaminase; ALP: alkaline phosphatase; TP: total protein; ALB: albumin.

Results on serum biochemistry showed that albumin and alkaline phosphate were significantly (P>0.05) affected by *Eupatorium odoratum* supplementation. Albumin values were statistically the same in treatment 1 (26.75±0.04g/l), treatment 3 (27.33±0.29g/l) and treatment 4 (26.82±0.41g/l) but higher than treatment 2 (28.18±0.09g/l). Alkaline phosphatase value was highest in treatments 3 and 4 (122.64±3.45 Iµ/L and 127.45±2.79 Iµ/L, respectively) followed by treatment 1 (102.40±5.10 Iµ/L) and least in treatment 2 (82.85±9.15 Iµ/L).

The aminotransferases, AST, and ALT, are typically inside the liver cells and are released into the blood when liver cells are damaged; increased levels are therefore indicative of liver damage (Okwesili et al., 2021). The findings of this study showed that supplementing *Eupatorium odoratum* leaf meal did not cause liver damage in the broilers. Total protein is a composite of albumin and globulin contents in the blood and are often used in diagnosing diseases and monitoring changes in the health status of farm animals (Oleforuh-Okoleh et al. 2015). Low levels of albumin could be

associated with presence of infection. The total protein of finisher birds in various treatments in this study were not influenced by *Eupatorium odoratum* inclusion. Moreover, albumin levels were improved among birds given 100g *Eupatorium odoratum*. Similar results were reported in the findings of Oleforuh-Okoleh et al. (2015) who fed *Vernonia amygdalina* leaf extract. ALP is a group of enzyme that initiates catalytic activity for degeneration of phosphate esters and separation of phosphoric acid molecules (Knits, 2008). Nonetheless, as an enzyme responsible for dephosphorylation of a substrate, it is produced in all types of tissues in the body, but it gets activated in alkaline pH. Therefore, elevated levels of ALP can be mostly seen in liver damages (Senanayake et al. 2015) at higher levels of *E. odoratum* above 200mg and may be attributed to longer duration of feeding (Oni et al. 2020).

Carcass Characteristics of Broiler Finisher Birds fed Diets Containing *Eupatorium Odoratum* Leaf Meal

Table 5: Carcass characteristics of broiler birds fed *Eupatorium odoratum* leaf meal

Parameters	Treatment 1 (control)	Treatment 2 (100mg <i>E.O</i>)	Treatment 3 (200mg <i>E.O</i>)	Treatment 4 (300mg <i>E.O</i>)	P- value
Live weight (g)	2566.01±1.21 ^b	2384.44±1.04 ^c	2577.24±1.50 ^a	2571.34±0.57	0.00**
	2053.35±1.47 ^b	1717.47±1.01 ^c	2143.81±1.12 ^a	2049.38±0.57 ^b	0.00**
De-feathered weight (g)	1500.59±1.50 ^b	1297.33±1.01 ^c	1582.84±1.02 ^a	1489.73±0.57 ^b	0.00**
Dressed weight (g)	58.22±0.02 ^b	54.41±0.02 ^c	61.68±0.02 ^a	57.94±0.57 ^b	0.00**
Dressing percentage (%)	32.93±1.21	32.86±1.04	33.00±1.50	32.95±0.57	1.00 ^{NS}
Breast cut (%)	23.95±1.31	23.30±1.03	23.72±1.05	23.51±0.57	0.98 ^{NS}
Back cut (%)	16.50±1.23	15.71±1.04	16.60±1.07	16.40±0.57	0.94 ^{NS}
Thigh (%)	16.54±1.21	15.59±1.02	16.70±1.50	16.55±0.57	0.89 ^{NS}
Drumstick (%)	0.26±0.06	0.63±0.23	0.49±0.35	0.52±0.11	0.69 ^{NS}
Kidney (%)	0.15±0.02	0.17±0.09	0.16±0.06	0.14±0.11	1.00 ^{NS}
Heart (%)	0.06±0.02	0.06±0.03	0.06±0.02	0.06±0.05	1.00 ^{NS}
Spleen (%)	1.46±0.06	1.78±0.09	1.52±0.12	1.50±0.11	0.25 ^{NS}
Liver (%)					

^{abc}: Means on the same row with different superscripts are significantly different (P<0.05).

The results on carcass characteristics of broiler birds fed *Eupatorium odoratum* leaf meal presented in Table 11 revealed that live weight, de-feathered weight, dressed weight, and dressing percentage were significantly (P<0.05) influenced by the treatment administered. The highest live weight was obtained from the birds on *Eupatorium odoratum* leaf meal supplemented at 200mg while the least weight was obtained from those on 100mg of *Eupatorium odoratum*. The same trend was observed for the de-feathered weight, dressed weight, and dressing %. Results on the relative percentage of breast cut, back cut, thigh, drumstick, kidney, heart, spleen and liver showed no significant (P>0.05) differences among the treatment groups.

Amounts of phyto-constituents in different concentrations and methods of extractions of meal could be suggested to be responsible for the change in the carcass characteristics of broilers (Dotas et al. 2014). Therefore, in relative to this study, most parameters were not significantly (P<0.05) affected by *Eupatorium odoratum* leaf meal diet which could be as a result of levels of inclusion or extraction method used when compared to the literature. The significant differences observed in dressed percentage is similar to the report of

Oloruntola et al. (2021) but not in agreement with the findings of Olumide and Akintola (2018) who reported similar values in birds of both control and treated groups. The variation in the reports could be attributed to the mode of administration of the *Eupatorium odoratum*. The former researchers used leaf extract while leaf meal was used in this present study. The similarity in the internal organs suggests the support of the extract to the normal functioning of the birds' internal organs. It has been reported by Ayodele et al. (2016) that the presence of toxic substances in the diet could lead to an increase or decrease in the relative weight of the birds internal organs. The improved live weight, dressed weight, and dressing percentage recorded in the birds fed *Eupatorium odoratum* leaf meal agreed with the previous reports that supplementation of the broiler chicken diets with phytochemicals improved the carcass weight and dressing percentage (Kanduri et al. 2013; Valenzuela-Grijalva et al. 2017). This proposes that the phyto-genic supplements used in this study have bioactive compounds that can modulate metabolic reactions in the animals' body system in a similar pattern with β -adrenergic agonist compound (Valenzuela-Grijalva et

al. 2017). On this note, one could say that *Eupatorium odoratum* could improve the performance and healthy state of broilers thereby standing as a growth promoter possessing both medicinal and nutritional qualities.

CONCLUSION AND RECOMMENDATION

The results of this study showed that *Eupatorium odoratum* leaf meal had several positive effects on the growth performance (final body weight, weight gain, feed conversion ratio and mortality reduction), haematology (packed cell volume, haemoglobin, white blood cells and monocytes) and carcass characteristics (live weight, de-feathered weight, dressed weight and dressing percentage) of broiler birds. Furthermore, serum biochemical constituent of albumin was also improved by *Eupatorium odoratum* leaf meal and the increment in alkaline phosphate which depicts the extent the body system of the birds reacted in the presence of anti-nutritional factors.

On this note, it is recommended that the use of *Eupatorium odoratum* leaf meal can improve broiler performance better at 200mg/kg diet for an efficient production and supply of animal protein. For the best result to be achieved, farmers are advised to properly process the leaf of *Eupatorium odoratum* properly in order to get rid of the anti-nutrient factors.

REFERENCES

- Abd El-Hack, ME and Alagawany M. (2015). Performance, egg quality, blood profile, immune function, and antioxidant enzyme activities in laying hens fed diets with thyme powder. *J. Anim. Feed Sci.* **24**: 127–133.
- Abdelli N, Solà-Oriol D and Pérez JF. (2021). Phytogetic Feed Additives in Poultry: Achievements, Prospective and Challenges. *Animals*. **11**: 3471.
- Aikpitanyi KU and Egweh NO. (2020). Haematological and biochemical profile of broiler chickens fed diets containing ginger and black pepper additives. *Nigerian J. Anim. Sci.* **22(2)**: 114-125.
- Akhter S, Hossain MW, Sultana S, Jharna JF, Meghla NS, Alam R and Rahman MM. (2022). *Ruellia prostrata* Poir. activity evaluated by phyto-constituents, antioxidant, anti-inflammatory, antibacterial activity, and in silico molecular functions. *Journal of Saudi Chemical Society*. **26(1)**: 101401.
- Alabi RA and Chime CC. (2008). *Impact of food production on food importation in Nigeria*. Final report presented to African Economic Research Consortium (AERC), Nairobi, Kenya.
- Alemu EA. (2020). Malnutrition and Its Implications on Food Security. In: Leal Filho, W., Azul AM, Brandli L, Özuyar PG, Wall T. (eds) Zero Hunger. Encyclopedia of the UN Sustainable Development Goals. Springer, Cham.
- Alisi CS, Onyeze GOC, Ojiako OA and Osuagwu CG. (2011). Evaluation of the Protective Potential of *Chromolaena odorata* Linn. Extract on Carbon Tetrachloride-Induced Oxidative Liver Damage. *International Journal of Biochemistry Research & Review*. **1(3)**: 69-81.
- Al-Khalaifah HS. (2018). Benefits of probiotics and/or prebiotics for antibiotic-reduced poultry. *Poult. Sci.* **97**: 3807–3815.
- Al-Nedawi A. (2018). Reference hematology for commercial Ross 308 broilers. *Online Journal of Veterinary Research*. **22(7)**: 566-570.
- Al-Yasiry ARM and Kiczorowska B. (2016). Frankincense – therapeutic properties. *Poet Hig MedDosw*. **70**: 380-391.
- Amaliah UN, Johannes E, Hassan MS and Tambaru E. (2019). The Use Extract of Siam Leaf *Eupatorium odoratum* L. as Alternative Material in Lowering Blood Glucose. *International Journal of Applied Biology*. **3(1)**: 15-23.
- Apori SO, Long RJ, Castro FB and Trskov ER. (2000). Chemical composition and nutritive value of leaves and stems of tropical weed *Chromolaena odorata*. *Journal of Grass and Forage Science*, **55**: 77-81.
- Ayodele SO, Oloruntola OD and Agbede JO. (2016). Effect of diet containing *Alchornea cordifolia* leaf meal on performance and digestibility of weaner rabbits. *World Rabbit Science*. **24**: 201–206.
- Bahman AH, Alireza, T and Siamak AR. (2011). Comparative Study on Blood Profiles of Indigenous and Ross-308 Broiler. *Breeders Global Veterinaria*. **7(3)**: 238-241.
- Bamikole MA, Ikhatua UJ and Osemwenkhae AE. (2004). Converting bush to meat: A case of *Chromolaena odorata* feeding to rabbits. *Pakistan Journal of Nutrition*. **3(4)**: 258-261.
- Breinholt KA, Gowen FA and Nwosu CC. (1981) Influence of environmental and animal factors day and night grazing activity of imported Holstein Friesian cows in the humid lowland tropics of Nigeria. *Trop Animal Produc.* **6**: 4-5.
- Dhabhar FS. (2000). Acute stress enhances while chronic stress suppresses skin immunity: the role of stress hormones and leukocyte trafficking. *Annals of the*

New York Academy of Sciences. **917(1)**: 876-893.

- Donkoh A, Atuahene C, Anang D, Badu-Botah EK and Boakye KT. (2002). Response of broiler chickens to the dietary inclusion of *Chromolaena odorata* leaf meal. *Journal of Animal and Feed Sciences*. **11(2)**: 309-319.
- Dotas V, Bampidis VA, Sinapis E, Hatzipanagiotou A and Papanikolaou K., (2014). Effect of dietary field pea (*Pisum sativum* L.) supplementation on growth performance, and carcass and meat quality of broiler chickens. *Livestock Science*. **164**: 135–143.
- Eguzoie B. (2019). *Nigeria's livestock production has to increase six-folds to satisfy animal protein demand*. Online article retrieved November 16th, 2021 from <https://www.businessamlive.com/nigerias-livestock-production-has-to-increase-six-folds-to-satisfy-animal-protein-demand-says-nias/>
- Ekwe OO, Uzoma CC, Ede VO and Nneji CP. (2017). Effect of *Eupatorium odoratum* and *Ocimum gratissimum* (Scent leaf) leaf extracts on growth, haematology and cost benefit of broiler chicken. *Asian J. Poult. Sci*. **11(2)**: 90-95.
- Escobedo del Bosque CI, Altmann BA, Ciulu M, Halle I, Jansen S, Nolte T and Mörlein D. (2020). Meat quality parameters and sensory properties of one high-performing and two local chicken breeds fed with *Vicia faba*. *Foods*. **9(8)**: 1052.
- FAO (2018). Africa Sustainable Livestock 2050: Livestock and livelihoods spotlight. NIGERIA. Cattle and Poultry Sectors. Available at: <http://www.fao.org/3/CA2149EN/ca2149en.pdf>. Accessed on the 15th January, 2022.
- Godde CM, Mason-D'Croz D, Mayberry DE, Thornton PK and Herrero M. (2021). Impacts of climate change on the livestock food supply chain; a review of the evidence. *Global Food Security*. **28**: 100488.
- Gopi S, Amalraj A, Varma K, Jude S, Reddy PB, Divya C. and Thomas, S. (2018). Turmeric nanofiber-encapsulated natural product formulation act as a phyto-genic feed additive—A study in broilers on growth performance, biochemical indices of blood, and *E. coli* in cecum. *International Journal of Polymeric Materials and Polymeric Biomaterials*. **67(9)**: 581-588.
- Jha PN, Mallik SK, Saxena A, Shahi N, Das P, Giri AK and Pandey PK. (2022). Leaf powder of *Eupatorium odoratum* Enhances non-specific immune response and resistance to *Aeromonas hydrophila* Infection in *Cyprinus carpio* (Linn. 1758). *Indian Journal of Animal Research*. **56(7)**: 880-886.
- Jiwuba PC, Ogbuewu IP and Nwachukwuguru K. (2018). Performance and economy of production of broilers fed Siam weed (*Chromolaena odorata*) leaf meal (SWLM). *Tropical Animal Health and Production*. An Official Publication of Enugu State University of Science & Technology ISSN: (Print) 2315-9650 ISSN: (Online) 2502-0524
- Kanduri AB, Munde VK, Khan MA, Thakur PN, Saxena MJ, Ravikanth K, Thakur A and Maini S. (2013). Study on the comparative efficacy of natural growth promoter (AV/AGP/10) with antibiotic supplements on overall growth performance and intestinal micrometry of broiler birds. *British Microbiology Research Journal*. **3**: 623–634.
- Knits M. (2008). Chicken intestinal alkaline phosphatase. *J. Gene. Physio*. **43**: 1149-1169.
- Machebe NS, Agbo CU and Onuaguluchi CC. (2011). Oral administration of *Gongronema latifolia* leaf meal: Implications on carcass and haematological profile of broiler finisher raised in the humid tropics. *African Journal of Biotechnology*. **10(30)**: 5800-5805.
- Marianne DLP, Elin YS, Neng FK and Rosnani N. (2014). Antidiabetic Activity of Leaf Ethanol Extract of *Chromolaena Odorata* (L.) R.M. King on Induced Male Mice with Alloxan Monohydrate. *Journal of Natural*. **14(1)**: 1–4.
- Mba C. (2021). #ChartoftheDay: Total Poultry Population by Agricultural Zones in Nigeria. An online article retrieved on February 12th, 2022 from <https://www.dataphyte.com/latest-reports/economy/chartoftheday-total-poultry-population-by-agricultural-zone-in-nigeria/>
- Megersa M. and Woldetsadik S. (2022). Ethnobotanical study of medicinal plants used by local communities of Damot Woyde district, Wolaita zone, southern Ethiopia. *Nusantara Bioscience*, **14(1)**:
- Nemeth N, Rudnak I, Ymeri P. and Fogarassy C. (2019). The role of cultural factors in sustainable food consumption—An investigation of the consumption habits among international students in Hungary. *Sustainability*. **11(11)**: 3052.
- Ofomata GEK. (1978). The Nsukka Environment Enugu, Nigeria: Fourth Dimension Publishers.
- Okosun SE. and Oyedeji E. (2016). Effect of feeding graded levels of *Moringa oleifera* leaf meal (MOLM) on the performance and blood biochemistry of broiler chicks. *Nigerian Journal of Agriculture and Forestry*. **8(6)**: 57-61.
- Okwesili OR, Ikele CM, Ikeh NE, Anizoba NW. and Osita CO. (2021). Oral administration of *Vernonia amygdalina* leaf extract: implications on performance of broiler finishers raised in derived savannah. *Agro-Science*. **20(2)**: 86-91.
- Oleforah–Okoleh VU, Ndofor–Foleng HM, Olorunlekes O. and Uguru JO. (2015). Evaluation of growth performance, Haematological and serum biochemical response of broiler chickens to aqueous extract of ginger and garlic. *Journal of Agricultural*

Science. **7(4)**: 167–73.

- Oloruntola OD, Adu OA, Gbore FA, Falowo AB. and Olarotimi OJ. (2021). Performance of broiler chicken fed diets supplemented with *Irvingia gabonensis* kernel powder and *Ocimum gratissimum* leaf powder. *Slovak Journal of Animal Science*. **54(1)**: 7–20.
- Olumide MD. and Akintola AS. (2018). Effect of scent leaf meal (*Ocimum gratissimum*) supplementation on performance, carcass and meat quality of broiler chicken. *Nigerian Journal of Animal Production*. **45(3)**: 228–236.
- Osho IB, Akindahunsi A, Igbasan FA, and Adekunle DJ. (2014). Effect of orally administered bitter leaf (*Vernonia amygdalina*) extract on the growth performance and haematological parameters of broiler chicken. *J. Vet. Med. Anim. Health*. **6**: 251–256.
- Ricke SC, Dittoe DK and Richardson KE. (2020). Formic acid as an antimicrobial for poultry production: A review. *Front. Vet. Sci*. **7**: 1–13.
- Schneitz C. and Hakkinen M. (2016). The efficacy of a commercial competitive exclusion product on *Campylobacter* colonization in broiler chickens in a 5-week pilot-scale study. *Poult. Sci*. **95**: 1125–1128.
- Senanayake SSH, JGS. Ranasinghe R. Waduge K. Nizanantha and PABD. Alexander. (2015). Changes in the Serum Enzyme Levels and Liver Lesions of Broiler Birds Reared Under Different Management Conditions. *Tropical Agricultural Research*, 26 (4): 584–595.
- Swain PS, Rao SBN, Rajendran D, Dominic G. and Selvaraju S. (2016). Nano zinc, an alternative to conventional zinc as animal feed supplement: A review. *Anim. Nutr*. **2**: 134–141.
- Tiamiyu AM. and Okunlade OA. (2021). Benefits and detriments of Siam weed (*Chromolaena odorata*): A review. *Biochemistry and Biotechnology Research*. **8(2)**: 21–28.
- Toghyani M, Shahryar HA. and Zamanizad M. (2011). Assessment of growth performance, immune responses, serum metabolites, and prevalence of leg weakness in broiler chicks submitted to early-age water restriction. *Tropical Animal Health and Production*. **43**: 1183–1189.
- Torres-Pitarch A, Manzanilla EG, Gardiner GE, O'Doherty JV. and Lawlor PG. (2019). Systematic review and meta-analysis of the effect of feed enzymes on growth and nutrient digestibility in grow-finisher pigs: Effect of enzyme type and cereal source. *Anim. Feed Sci. Technol*. **251**: 153–165.
- Ukam VE, Agwunobi LN. and Oko OOK. (2020). Carcass and organ characteristics of broiler finisher chickens fed matured sun-dried nypa palm fruit (*Nypa fruticans*) nut meal. *Nigerian J. Anim. Sci*. **22(3)**: 158–164.
- United States Department of Agriculture – USDA (2018). Chickens and Eggs. ISSN: 1948-9064 Available at: https://www.nass.usda.gov/Publications/Todays_Reports/reports/ckeg1218.pdf.
- Valenzuela-Grijalva NV, Pinelli-Saavedra AP, MuhliaAlmazan A, Dominguez-Diaz D. and Gonzalez-Rios H. (2017). Dietary inclusion effects of phytochemicals as growth promoters in animal production. *Journal of Animal Science and Technology*. **59(8)**: 1–17.
- Vastolo A, Calabrò S. and Cutrignelli MI. (2022). A review on the use of agro-industrial CO-products in animals' diets. *Italian Journal of Animal Science*. **21(1)**: 577–594.
- Wong JT, de Bruyn J, Bagnol B, Grieve H, Li M, Pym Rz, and Alders RG. (2017). Small-scale poultry and food security in resource-poor settings: A review. *Global Food Security*. **15**: 43–52.
- Zhai H, Liu H, Wang S, Wu J. and Kluentner AM. (2018). Potential of essential oils for poultry and pigs. *Anim. Nutr*. **4**: 179–186.