

Table 1.Positivity rates of study groups according to epithelial, stromal and adipose tissue

		Epithelial		Stromal		Adipose tissue		<i>Total</i>		p
		n	%	n	%	n	%	<i>n</i>	<i>%</i>	
SHAM	Negative	4	80.0	2	40.0	2	40.0	8	53.3	0.322 ^{b1}
	Positive	1	20.0	3	60.0	3	60.0	7	46.7	
TOES 14	Negative	6	75.0	7	87.5	7	87.5	20	83.3	0.751 ^{b1}
	Positive	2	25.0	1	12.5	1	12.5	4	16.7	
TOES 28	Negative	6	75.0	8	100	5	62.5	19	79.2	0.083 ^{b1}
	Positive	2	25.0	0	0.0	3	37.5	5	20.8	
p		0.973 ^{b2}		0.021^{b2}		0.184 ^{b2}		0.090 ^{a3}		

^a: Pearson Chi-square

^b: Likelihood ratio Chi-square

Table 2.Positivity rates of study groups according to epithelial, stromal and adipose tissue

		Epithelial		Stromal		Adipose tissue		<i>Total</i>		p
		n	%	n	%	n	%	<i>n</i>	<i>%</i>	
SHAM	Negative	4	80.0	2	40.0	2	40.0	8	53.3	0.322 ^{b1}
	Positive	1	20.0	3	60.0	3	60.0	7	46.7	
TOES 14	Negative	6	75.0	7	87.5	7	87.5	20	83.3	0.751 ^{b1}
	Positive	2	25.0	1	12.5	1	12.5	4	16.7	
TOES 28	Negative	6	75.0	8	100	5	62.5	19	79.2	0.083 ^{b1}
	Positive	2	25.0	0	0.0	3	37.5	5	20.8	
p		0.973 ^{b2}		0.021 ^{b2}		0.184 ^{b2}		0.090 ^{a3}		

^a: Pearson Chi-square; ^b: Likelihood ratio Chi-square

¹: Regions (Epithelial, Stroma and Adipose Tissue) were compared in each group

2: Groups (SHAM, TOES 14 and TOES 28) compared in each region

3: Groups were compared without considering the regions (Epithelial, Stroma, Adipose Tissue)

Graph-1: Graphical evaluation of all results







