

EFFECT OF FERMENTATION PERIODS ON SOME ATTRIBUTES OF SORGHUM CHINCHIN

ABSTRACT

Fermentation is a cheap method of food preservation and processing. Cereals can be fermented to provide commercially available snacks, meals, and alcoholic beverages. This research examined at how the length of the fermenting time affects some sorghum chinchin quality attributes (in hours). The sorghum grains were subjected to natural fermentation for 0, 12, 24, 36, 48, and 60 hours. They were then dried and processed into flour, and the dough was fried in oil to form Chinchin. Analyses of proximate composition, colour, texture profile, functional and sensory properties were carried out on the samples. The data obtained were further analyzed using ANOVA. Analysis of proximate composition showed an increase in the fat, crude fibre and crude protein contents but a decrease in the carbohydrate content as the fermentation period progresses; $19.12\pm 0.01\%$ to $19.44\pm 0.02\%$, $3.27\pm 0.01\%$ to $4.4\pm 0.00\%$, $3.85\pm 0.01\%$ to $4.19\pm 0.01\%$, $64.48\pm 0.01\%$ to $61.05\pm 0.01\%$ respectively, while the values obtained from the control were $17.33\pm 0.01\%$, $3.31\pm 0.01\%$, $3.72\pm 0.01\%$, $65.68\pm 0.08\%$ respectively. Oil uptake for the control sample was $18.64\pm 0.69\%$ and a decrease from $17.71\pm 0.03\%$ to $12.42\pm 0.08\%$ was noted for the fermented samples as the fermentation period increased. 36-hour fermented sample had the lowest value at 7.0 ± 1.08 , whereas the control samples and the 60-hour fermented sample had the highest overall acceptability. Implementing the consumption of snacks produced from 100% fermented sorghum as part of the diet can help prevent diet-related diseases and improve nutrition. Also, the commercial production of the snacks could provide additional socioeconomic benefits and increase the economic value of sorghum.

Subjects: Environment & Agriculture; Bioscience; Food Science & Technology

Keywords: Sorghum; Wheat; Fermentation; Chinchin; Proximate; Sensory Evaluation

1. INTRODUCTION

Sorghum is the fifth most important cereal crop in the world after rice, wheat, corn and barley. It is the main cereal food for over 750 million people living in semi-arid tropical regions of Africa, Asia and Latin America [1]. Sorghum locally called guinea corn and “*dawa*” in Nigeria is a gluten-free grain that has the potential to be used as an alternative to wheat flour [2]. Sorghum grain is a rich source of various phytochemicals such as tannins, phenolic acids etc., which have health-promoting activities, anti-cancer, anti-tumor, antidiabetic and anti-obesity properties [3]. Sorghum grain is a rich source of macronutrients (carbohydrates, proteins, and fat) and micronutrients (mineral and vitamins). It is rich in magnesium, iron, manganese and phosphorus [4]. Sorghum grains contain resistant starch, which makes it interesting for obese and diabetic people, as the digestibility of whole sorghum is slower than other major cereals, leading the slow release of glucose into the blood. Therefore, the energy released is fully utilized and prevents accumulation of fat. It is also recommended as gluten-free food for celiac patients [5]. Although cereal grains constitute a major source of dietary nutrients worldwide, they are deficient in some basic components (e.g. Essential amino acids). Fermentation may be the simplest and most economical way of improving their nutritional value and functional qualities. [6].

Fermentation is well-known to improve the sensory properties of food by imparting unique flavours, textures and aromas [7]. It is also used to improve the bioavailability and bio-accessibility of nutrients, reduce antinutritional factors (such as lectins, phytic acid, proteinase inhibitors, oxalic and tannins acids) and pathogenic microorganisms, preserve food products as well as to enhance the economic value [8]. As a result, fermented foods have become a very important part of the human diet worldwide. Fermented food products have been in existence since the arrival of human civilization and are likely to be with us far into the future. Fermentation is, thus, an age-long food processing technique used to transform food products [9,10], with different food crops (cereals, legumes, as well as fruits and vegetables) used as starting raw materials.

Increasing fermentation period results in a decrease in tannin content, starch, amylopectin, and water and oil absorption and an increase in the value of whiteness, dietary fiber, amylose, and viscosity of sorghum flour [11]. Also, in a study done by [12], It was noted that increased fermentation time and temperature led to an increase in mean levels of some phenolic compounds of Ting obtained from whole grain sorghum.

Studies have shown that snacks can be used to increase the nutritional status of consumers by incorporating nutrients such as protein and fibre from plant sources with significant health benefits [13]. Use of sorghum grains as composite flour have previously been studied

in the production of bread, extruded snacks, biscuits among others. But, the utilization of 100% sorghum flour in food production and its acceptability is not yet widely examined and information on its use as snacks is insufficient.

Chin-chin is a fried snack popular in West African countries especially Nigeria. It is a sweet hard doughnut-like fried product which is sometimes a baked dough of wheat flour, with eggs and other ingredients [14]. The flour is mixed to form an elastic dough which is properly kneaded, rolled and cut into desired shapes. The shaped flat dough is then deep fried in hot oil until slightly golden brown then scooped out to let the oil drain [15].

The objective of this research work is to evaluate the effect of fermentation periods on the proximate composition, sensory characteristics, moisture loss, oil uptake and textural properties of a chinchin snack produced from 100 % sorghum flour using 100% wheat chinchin as control.

2. MATERIALS AND METHODS

2.1 MATERIALS

Materials used (*Sorghum bicolor* grains, margarine, sugar, nutmeg, salt and baking powder) were purchased at Osiele Market, Abeokuta. The production was carried out in the food processing laboratory in the Department of Food Science and Technology, Federal University of Agriculture Abeokuta, Ogun State, Nigeria.

2.2. Fermentation of sorghum grains

1.3kg of whole sorghum grains were sorted manually to remove stones, damaged grains and foreign particles. The grains were put in a plastic container, steeped in 1 litre of distilled water and covered tightly. The grains were subsequently subjected to natural fermentation at 37°C. This was repeated separately for different periods of time (12hr, 24hr, 36hr, 48hr, and 60hr).

250g of unfermented whole sorghum grain was also sorted manually to remove damaged grains and all foreign particles. The grains were rinsed thoroughly with distilled water without allowing fermentation to take place (0hr).

Afterwards, the unfermented (0hr) and fermented grains (12hr, 24hr, 36hr, 48hr, and 60hr) were dried at 70°C for 4hrs in hot air oven, exposed to air to cool and the samples were labelled according to the fermentation period.

2.3 Sorghum flour production

The dried, cooled fermented grains was milled using a single flour milling machine into flour (to be used in the chinchin production). The flour was sieved, packaged in airtight polyethylene bags and stored at 37°C in covered plastic buckets for storage (4 days before the chinchin snacks production).

2.4 Chinchin preparation

100g of fermented sorghum flour was poured in a bowl, then mixed with 30g of sugar, 20g of margarine, 1.42g of baking powder, 2.7g of salt and 1g of nutmeg. 100g of fermented flour was poured in another bowl and a well was made at the center; 118ml of water was heated to a temperature of 32°C and poured in the center; the flour was gently folded in with a turning stick 3 times and poured immediately to the first flour mixture. The lumps formed were separated with fingertips till all ingredients were well combined and resembled breadcrumbs. Eggs whisked until light, was added to the mixture, mixed with finger tips into smooth dough leaving the side of the bowl clean. It was rolled evenly on a floured board and cut into desired shape and size; length and thickness. Deep-fried in oil until attractively brown. Excess oil was drained. The product was then cooled and packaged in high density polyethylene bag for storage until the chinchin was evaluated. This was repeated for all samples.

2.4.1 Proximate composition

Protein, fat, crude fibre, moisture and ash were determined by the methods of Analysis of the Association of Official Analytical Chemists [16], while carbohydrate was determined by difference.

2.4.2. Functional properties of the chinchin

Moisture loss

Moisture content of the chinchin was determined using methods of Analysis of the Association of Official Analytical Chemists [16].

MOISTURE LOSS: (Moisture content of fried chinchin) - (Moisture content of the dough).

Oil uptake

Oil content of the chinchin was determined using methods of Analysis of the Association of Official Analytical Chemists [17] method

OIL UPTAKE: (Oil content of fried chinchin) – (Oil content of dough)

2.4.3 Colour analysis

Each sample of the chinchin was analyzed using a colorimeter to determine their brightness, redness and yellowness by the application of Beer-Lambert's law.

2.4.4 Texture Analysis

The instrumental texture measurements of the sorghum chinchin was analyzed with a TA.XT.plus Texture Analyzer (Stable Microsystems, Godalming, UK) provided with Texture expert software. The chinchin (with a thickness of 2mm) was placed horizontally directly under the probe. A double compression test (texture profile analysis) was performed with a 75 mm diameter flat-ended cylindrical probe (P/75) and compression to 50% of the initial height at a speed of 1 mm/s with 5s waiting time between the two cycles.

2.4.5 Sensory evaluation

The sensory attributes of the *chinchin* was determined by using simple hedonic tests as described by [18]. This was done using a 20-member panel comprising of students of the department of Food science and technology, Federal University of Agriculture, Abeokuta, who were familiar with the sensory attributes of chin-chin. Each panelist was asked to score each attribute on a 9-point hedonic scale where 1 and 9 represent dislike extremely and like extremely, respectively. The attributes that were evaluated includes appearance, colour, taste, texture, aroma and overall acceptability.

2.4.6 Data analysis

All data obtained was subjected to analysis of variance (ANOVA). Means was separated using Duncan's multiple range test (DMRT) using the Statistical package for social sciences (SPSS) version 21.0 (SPSS Inc., Chicago, IL).

3. RESULT AND DISCUSSION

3.1 Proximate composition of the fermented sorghum chinchin sample

The proximate composition of the chinchin samples is shown in Table I. The moisture content increased from 6.22 to 6.37 % as the fermentation period increased and was significantly different ($p < 0.05$) from each other. The values were within the range reported to have no adverse effect on quality attribute of the product [13]. The low moisture content could reduce the growth of micro-organism thereby increasing the shelf life of the product.

Dietary fat functions in the increase of palatability of food by absorbing and retaining flavours [19]. The fat content increased from 19.12 to 19.44 % with a significant difference ($p < 0.05$) from each other, There was an increase in the fat content as the fermentation period is increased due to an increase in the activity of the lipolytic enzymes in hydrolyzing fat to glycerol and fatty acids [20].

Ash content is an indication of the mineral content of a food [21]. The ash content ranged between 3.07 to 4.59 % with 48hr having the highest. The ash content has been reported to increase with fermentation according to [22] and [23].

The values for crude fibre ranged between 3.12 to 4.40 % with 60hr having the highest while 12hr had the least. Crude fibre content increases with an increase in the fermentation period. Sorghum contains high level of fibre and this was further increased by fermentation [20]. Crude fibre helps in the prevention of heart diseases, colon cancer and diabetes [24].

The protein value ranged from 3.85 to 4.19%, 60hr had the highest and 0hr had the lowest value with a significant difference ($p < 0.05$) for all samples. The protein content increased with an increase in the fermentation period, which could be due to synthesis of protein by micro-organisms. Fermentation increases protein value [25]. Protein is needed for growth, maintenance and regulation of the body process.

The carbohydrate content of the chinchin ranged between 64.48 to 61.05 % with a significant difference ($p < 0.05$). The carbohydrate value decreased as fermentation period increased. This agrees with [26], who reported that carbohydrate content was reduced with fermentation due to the increase of alpha-amylase activity which hydrolyses starch to simple sugars.

Table I: Proximate analysis for the fermented sorghum chinchin sample

Fermentation Period (hrs)	Moisture Content (%)	Fat (%)	Ash (%)	Crude Fibre (%)	Crude Protein (%)	Carbohydrate (%)
Control	6.46±0.02^e	17.33±0.01^a	3.52±0.01^d	3.31±0.01^c	3.72±0.01^a	65.68±0.08^g
0	6.22±0.01 ^d	19.12±0.01 ^f	3.07±0.01 ^f	3.27±0.01 ^g	3.85±0.01 ^d	64.48±0.01 ^a
12	6.24±0.01 ^a	19.15±0.01 ^b	3.11±0.01 ^a	3.12±0.01 ^b	3.85±0.01 ^b	64.57±0.00 ^e
24	6.31±0.01 ^a	19.17±0.01 ^{bc}	3.48±0.01 ^b	3.51±0.01 ^a	4.09±0.01 ^b	63.46±0.02 ^f
36	6.34±0.01 ^b	19.31±0.01 ^c	3.98±0.01 ^c	4.08±0.01 ^d	4.11±0.01 ^c	62.2±0.05 ^d
48	6.36±0.01 ^c	19.36±0.01 ^d	4.59±0.01 ^e	4.32±0.01 ^e	4.17±0.01 ^c	61.22±0.02 ^c
60	6.37±0.01 ^{cd}	19.44±0.02 ^e	4.56±0.01 ^f	4.4±0.00 ^f	4.19±0.01 ^d	61.05±0.01 ^b

Mean values with different superscripts within the same column are significantly different (p<0.05)

Control: 100% wheat flour

3.2 Functional properties of Fermented Sorghum Chinchin (FSC)

3.2.1 Moisture loss for the Fermented Sorghum Chinchin (FSC)

The functional properties are those parameters that determines the application and end use of food materials for various food products. Moisture loss plays a determining role in oil uptake due to its ability to create spores [27]. Moisture loss is concerned with the difference between the Moisture content of the dough and Moisture content of the fried chinchin.

Evaluation of the Moisture loss for each of the Sample was carried out as shown in Table II. The values for the Moisture content of the dough ranged from 31.44 to 26.72 with 24hrs having the highest value and 60hrs having the lowest value. There was a significant difference of (p<0.05). Data showed that there was a decrease in moisture content of the dough which is due to decrease in water binding capacity of sorghum flour during fermentation [28]. The moisture content of the fried snack was in the range of 12.77 to 10.21 with 12hrs having the highest and 60hrs having the lowest. There was significant difference of (p<0.05) between the values. The value of moisture loss was between 18.93 and 15.75 with 24hrs having the highest and 0hrs having the lowest. The data significantly differed with (p<0.05).

Table II: Moisture loss for the Fermented Sorghum Chinchin (FSC)

Fermentation Period(hrs)	Dough Moisture content (%)	Snack Moisture content (%)	Moisture loss (%)
Control	25.68±0.01^a	10.21±0.01^a	15.47±0.02^a
0	28.32±0.01 ^c	12.57±0.00 ^b	15.75±0.01 ^b
12	29.93±0.01 ^e	12.77±0.01 ^d	17.17±0.02 ^d
24	31.44±0.02 ^g	12.51±0.01 ^f	18.93±0.03 ^f
36	30.11±0.01 ^f	12.38±0.01 ^e	17.74±0.02 ^e
48	28.86±0.00 ^d	12.34±0.01 ^c	16.53±0.01 ^c
60	26.72±0.01 ^b	10.21±0.01 ^c	16.51±0.02 ^c

Mean values with different superscripts within the same column are significantly different (p <0.05)

Control: 100% wheat flour

3.2.2 Oil uptake for the Fermented Sorghum Chinchin (FSC)

Evaluation of the oil uptake for each of the Sample was carried out as shown in Table III. Oil absorption capacity is important since oil acts as flavor retainer and increases mouth feel of foods [29].

The values for the oil content of the fried snack ranged from 21.14 to 16.28 with 0hrs having the highest value and 60hrs having the lowest values. The values were significantly different from each other (p<0.05). The values decreased with increase in fermentation period.

The values for the oil content of the dough ranged from 6.43 to 3.87 with 0hrs having the highest value and 60hrs having the lowest values. The values were significantly different from each other (p<0.05).

The values for the oil uptake of the fried snack ranged from 17.71 to 12.42 with 0hrs having the highest value and 60hrs having the lowest values. The values were significantly different from each other ($p < 0.05$). In line with [28], the values of the oil uptake by the fried snack decreased with an increase in fermentation period which could be as a result of decrease in water binding capacity of sorghum flour during fermentation.

Table III: Oil uptake for the Fermented Sorghum Chinchin (FSC)

Fermentation Period(hrs.)	Oil Content Fried (%)	Oil Content Dough (%)	Oil uptake (%)
Control	25.15±0.67^e	6.51±0.01^g	18.64±0.69^f
0	24.14±0.04 ^e	6.43±0.01 ^f	17.71±0.03 ^{ef}
12	22.65±0.04 ^d	5.19±0.01 ^e	17.46±0.06 ^d
24	21.02±0.94 ^c	5.05±0.04 ^d	15.97±0.98 ^c
36	20.14±0.04 ^c	4.92±0.02 ^c	15.22±0.01 ^c
48	18.24±0.01 ^b	4.66±0.06 ^b	13.58±0.07 ^b
60	16.28±0.04 ^a	3.87±0.04 ^a	12.42±0.08 ^a

Mean values with different superscripts within the same column are significantly different ($p < 0.05$)

Control: 100% wheat flour

3.3 Colour analysis for Fermented Sorghum Chinchin (FSC)

Colour is an important aspect in the market. Consumer perceptions about some products are based on color and many foods are associated with a specific color. The surface color values of the chinchin were analyzed using a CIE lab color system L value. According to the L, a, b type of scales: L (lightness) axis—0 is black, 100 is white; a (red-green) axis—positive values are red; negative values are green and 0 is neutral; and b (yellow-blue) axis—positive values are yellow; negative values are blue and 0 is neutral. All colors that can be perceived visually can be measured in any L, a, b scale. Lightness (L^*), a critical colour parameter of fried foods, is usually used as a quality control determinant.

Since lightness is a very important colour quality parameter, lower frying temperatures with lower boiling point of water are preferable to preserve the lightness and hence the attractiveness of fried products and so its adequate control is of utmost importance [30]. Table IV shows the result of the colour analysis done for the chinchin samples.

For all the frying treatments in this study, L^* values ranged from 43.12 to 34.64%. 0hr had the highest and 60hrs had the lowest value. However, there was a significant difference between all the samples at ($p < 0.05$) level of significance. There was a decrease in the lightness of the chinchin samples as a result of loss of lightness as the fermentation period increased.

The a^* values of the chinchin ranged from 9.94 to 9.19; 0hrs had the highest and 60hrs had the lowest value and there was a significant difference between all the samples at ($p < 0.05$) level of significance. This shows that there is a decrease in the redness of the chinchin sample as fermentation is increased.

The b^* values ranged from 21.33 to 17.11 with 0hrs having the highest and 60hrs having the lowest value and there was a significant difference between all the samples at ($p < 0.05$) level of significance as fermentation is increased.

The colour values (a/b) ranged from 0.55 - 0.69 with 0hrs having the lowest value and 60hrs having the highest value and there was a significant difference between all the samples at ($p < 0.05$) level of significance and this shows that there is an decrease in the colour value with increased fermentation period and this is similar to the observation made by [31] in which a darker colour was observed in the bread made by fermented sorghum flour. The dark colour could be as a result of higher reducing sugar content which undergoes non-enzymatic browning and the frying process. This showed that reducing sugars increased during fermentation as observed by [32].

Table IV: Colour analysis for Fermented Sorghum Chinchin (FSC)

Fermentation Period (hr)	L*	a*	b*	(a/b)
Control	45.83±0.35^d	14.43±0.16^c	31.13±0.28^e	0.67±0.01^c
0	43.12±0.37 ^c	9.94±0.14 ^b	24.3±0.35 ^d	0.55±0.01 ^a
12	38.4±0.05 ^b	9.3±0.00 ^a	20.15±0.01 ^c	0.60±0.0 ^b
24	37.40±0.2 ^b	9.33±0.06 ^a	17.89±0.44 ^b	0.67±0.01 ^c
36	37.29±1.04 ^b	9.33±0.12 ^a	17.16±0.06 ^a	0.68±0.00 ^c
48	37.11±1.67 ^b	9.15±0.01 ^a	17.18±0.04 ^a	0.60±0.0 ^b
60	34.64±0.45 ^a	9.19±0.08 ^a	17.11±0.08 ^a	0.69±0.01 ^c

Mean values with different superscripts within the same column are significantly different (p<0.05)

Control: 100% wheat flour

3.4 Texture profile analysis of Fermented Sorghum Chinchin (FSC)

Texture analysis is primarily concerned with measurement of the mechanical properties of a product, often a food product, as they relate to its sensory properties detected by human while applying controlled forces to the product and recording its response in the form of force, deformation and time. Texture measurements can be very valuable for the quality control and process optimization as well as for the development of new products with desirable properties and characteristics [33].

The mechanical characteristics parameters are the hardness, springiness, adhesiveness and cohesiveness [34]. Texture analysis is an objective physical examination of baked products and gives explicit information on the product's quality [35].

Table V shows the textural parameters assessed from texture profile analysis (TPA) test curves results of the chin-chin samples. Although hardness in snacks such as chin-chin is dependent on moisture content, moisture migration and redistribution of water, gluten-starch interactions [36], the fermentation period has no much effect on the hardness of the chinchin samples. 0hrs has the lowest value for hardness 36hrs has the highest values for hardness.

At the 48hr fermentation period, lowest value for springiness was recorded as shown in table V, this could be as a result of absence of gluten in the sorghum flour. Springiness refers to the rate at which a deformed product goes back to its un-deformed state, when the deforming force is removed while chewiness is the energy needed to masticate solid food to a state of readiness for swallowing [37] and is directly related to hardness and chewiness.

Table V: Texture profile analysis (TPA) of Fermented Sorghum Chinchin (FSC)

Fermentation Period (hr.)	Hardness (N)	Springiness	Adhesiveness (N.s)	Cohesiveness	Chewiness (N)	Fracturability (N)	Gumminess (N)	Stringiness (mm)
Control	163.55±0.21 ^a	0.02±0.00 ^a	0.02±0.01 ^a	0.04±0.01 ^a	0.08±0.00 ^a	23.45±0.04 ^a	7.22±0.01 ^a	4.53±0.00 ^f
0	297.4±412.25 ^{ab}	0.12±0.00 ^d	2.28±0.06 ^c	0.13±0.01 ^d	3.82±0.00 ^f	34.64±0.01 ^c	34.22±0.01 ^e	4.37±0.01 ^e
12	477.49±1.03 ^{ab}	0.12±0.00 ^d	0.03±0.00 ^a	0.05±0.00 ^b	3.49±0.01 ^e	31.04±0.01 ^b	31.66±0.04 ^d	4.37±0.01 ^e
24	585.65±0.07 ^b	0.08±0.01 ^c	0.23±0.01 ^b	0.05±0.00 ^b	1.12±0.00 ^c	50.83±0.01 ^e	31.38±0.06 ^c	3.58±0.01 ^a
36	585.91±0.34 ^b	0.09±0.00 ^c	0.01±0.00 ^a	0.16±0.00 ^e	2.16±0.00 ^d	57.51±0.11 ^d	46.14±0.14 ^g	3.83±0.00 ^c
48	332.2±0.46 ^{ab}	0.06±0.00 ^b	0.01±0.00 ^a	0.08±0.00 ^c	0.84±0.00 ^b	41.9±0.00 ^e	23.47±0.07 ^b	3.76±0.01 ^b
60	566.85±0.92 ^b	0.08±0.00 ^c	0.02±0.00 ^a	0.15±0.00 ^e	3.82±0.01 ^f	76.4±0.47 ^g	44.64±0.01 ^f	3.92±0.00 ^d

Mean values with different superscripts within the same column are significantly different (p<0.05). Values are means and standard deviations of two determination

Control: 100% wheat flour

3.5 Sensory characteristic of Fermented Sorghum Chinchin (FSC)

The result of the sensory evaluation of sorghum chin-chin are shown in Table VI. The colour, taste, texture, texture, aroma and overall acceptability of the composite chin-chin were not statistically different ($p>0.05$).

The result showed that the snack was liked for all the levels. In terms of colour, the values ranged from 7.65 to 6.25 with the control sample having the highest score and 0hr having the lowest. In terms of texture, the values ranged between 7.20 and 6.40 with 48hrs having the highest score and 0hrs having the least score. The high score rating for 48hrs could be as a result of the hardness due to increased fibre content due to fermentation which provides a crunchy sensation during mastication.

The texture was not statistically different ($p>0.05$) for all the samples. In a similar way, in terms of aroma, the mean values ranged from 6.9 to 6.40, with 24hrs having the highest and 48hrs having least score.

The high sensory score observed for aroma may be attributed to the development of flavour cause by the fermentation. No Significant difference ($p>0.05$) was observed for the samples. Taste ranged from 7.60 and 6.80 with 0hr having the highest while 24hr chinchin had the least. Sample for the control was the most acceptable followed by 60hrs but 36hrs chin-chin was the least accepted. Generally, the overall acceptability increased with fermentation.

Table VI: sensory analysis for the Fermented Sorghum Chinchin (FSC)

Fermentation Period(%)	Colour	Texture	Aroma	Taste	Overall Acceptability
Control	7.65±1.27 ^b	7.1±1.07 ^a	6.75±1.12 ^a	7.0±1.21 ^a	7.45±1.00 ^a
0	6.25±1.37 ^a	6.4±1.31 ^a	6.65±1.53 ^a	7.6±1.43 ^a	7.15±1.42 ^a
12	6.65±0.88 ^a	6.55±1.1 ^a	6.55±1.1 ^a	6.9±1.33 ^a	7.25±0.91 ^a
24	6.3±1.53 ^a	6.6±1.43 ^a	6.9±1.33 ^a	6.8±1.32 ^a	6.95±1.23 ^a
36	6.35±1.18 ^a	6.65±0.93 ^a	6.35±1.09 ^a	7.15±1.18 ^a	7.0±1.08 ^a
48	6.85±1.23 ^a	7.2±1.15 ^a	6.4±1.05 ^a	7.35±1.5 ^a	7.2±1.15 ^a
60	6.5±1.15 ^a	6.9±1.37 ^a	6.85±1.27 ^a	6.95±1.28 ^a	7.40±1.00 ^a

Mean values with different superscripts within the same column are significantly different ($p < 0.05$)

Control: 100% wheat flour

4. CONCLUSION

This study examined how some sorghum chinchin properties were affected by the length of the fermentation process. The results obtained showed a notable increase in the protein content, fat content and dietary fibre while the carbohydrate content decreased with the fermentation period. Also, a high overall sensory acceptability was recorded for the fermented sorghum snack.

As the incidence of celiac disease rises, gluten-free cereals like sorghum are being investigated as alternatives to wheat. Snacks are very popular in Nigeria and are consumed by majority of the population. Consumption of Fermented Sorghum Chinchin (FSC) should be promoted as a healthy approach to boost nutrition, improve digestion, and lower dietary-related problems like celiac disease, obesity, and diabetes. In addition, fermentation is a cost-effective technology and this innovative product will provide prospective excellent economic benefits and will greatly improve the economic value of sorghum.

Further research work on the microbial status and anti-nutritional properties of fermented sorghum chinchin is recommended. Also, storage studies should be conducted on the finished product to determine the shelf stability of the snack.

ACKNOWLEDGMENT

Authors will like to acknowledge the contribution of the Technical staff in the Department of Food Science and Technology, Federal University of Agriculture, Abeokuta, Nigeria.

FUNDING

The authors received no direct funding for this research.

COMPETING INTERESTS

The authors declare no competing interest.

REFERENCES

- Joint FAO/WHO2011 Food Standards Codex Committee on Contaminants in Foods (CCCF). Working paper on mycotoxins in sorghum.
- Li, X.R.; Ma, E. B.; Yan, L. Z.; Meng, H.; Du, X.W.; Zhang, S. W.; and Quan, Z. X. (2011) Bacterial and fungal diversity in the traditional Chinese liquor fermentation process. *Int. J. Food Microbiol.*, 146, 31–37.
- Awika J. M, and Rooney L. W. (2004) Sorghum Phytochemicals and their potential impact on human health. *Phytochemistry*. 65(9):1199-1221.
- Rajesh Devisetti, Sreerama N. Yadahally, and Sila Bhattacharya, (2014) Nutrients and antinutrients in foxtail and proso millet milled fractions: Evaluation of their flour functionality, *LWT - Food Science and Technology*, Volume 59, Issue 2, Part 1, Pages 889-895, ISSN 0023-6438, <https://doi.org/10.1016/j.lwt.2014.07.003>.
- Dicko MH, Gruppen H, Traoré AS, Voragen AG and van Berkel WJ (2006). Review: sorghum grain as human food in Africa: relevance of starch content and amylase activities. *African Journal of Biotechnology*, 5, 384-395.
- Taylor, J.R N., Barrlon, S. C, and Rooney, L.W. (2010) Pearl Millet – New developments in an ancient food grain. *Cereal Foods World* 55:16.
- Chaves-Lopez, C.; Serio, A.; Grande-Tovar, C.D.; Cuervo-Mulet, R.; Delgado-Ospina, J.; and Paparella, A. (2014) Traditional fermented foods and beverages from a microbiological and nutritional perspective: *The Colombian Heritage.Compr. Rev. Food Sci. Food Saf.* 13, 1031–1048.
- Hlangwani, E.; Adebisi, J.A.; Doorsamy, W.; and Adebo, O.A. (2020) Processing, characteristics and composition of umqombothi (a SouthAfrican traditional beer) Processes.,8, 1451.
- Adebisi, J.A.; Obadina, A.O.; Adebo, O.A.; and Kayitesi, E. (2018) Fermented and malted millet products in Africa: Expedition from traditional/ethnic foods to industrial value-added products. *Crit. Rev. Food Sci. Nutr.*,58, 463–474.
- Adebo, O.A. (2020) African sorghum-based fermented foods: Past, current and future prospects. *Nutrients* 12, 1111.
- Oluwafemi Ayodeji Adebo, Patrick Berka Njobeh, Janet Adeyinka Adebisi, and Eugenie Kayitesi, (2018) Co-influence of fermentation time and temperature on physicochemical properties, bioactive components and microstructure of ting (a Southern African food) from whole grain sorghum, *Food Bioscience*, Volume 25, Pages 118-127, ISSN 2212-4292, <https://doi.org/10.1016/j.fbio.2018.08.007>.
- Taiwo O. Osungbaro, (1990) Effect of fermentation period on amylose content and textural characteristics of Ogi (a fermented maize porridge), *Journal of Fermentation and Bioengineering*, Volume 70, Issue 1, Pages 22-25, ISSN 0922-338X, [https://doi.org/10.1016/0922-338X\(90\)90024-Q](https://doi.org/10.1016/0922-338X(90)90024-Q).
- Zazueta-Morales, J. J., Martínez-Bustos, F., JacoboValenzuela, N., Ordorica-Falomir, C., & Paredes-López, O. (2001). Effect of the addition of calcium hydroxide on some characteristics of extruded products from blue maize (*Zea mays* L.) using response surface methodology. *Journal of the Science of Food and Agriculture*, 81, 1379–1386. <http://dx.doi.org/10.1002/jsfa.951>

- Akubor, P. I. (2004). Protein contents, physical and sensory properties of Nigerian snack foods (cake, chin-chin and puff-puff) prepared from cowpea-wheat flour blends. *International Journal of Food Science and Technology*, 39, 419–424. <http://dx.doi.org/10.1111/ifs.2004.39.issue-4>
- Mepba, H. D., Achinewhu, S. N., & Aso, S. N. (2007). Microbiological quality of selected street foods in Port Harcourt, Nigeria. *Journal of Food Safety*, 27, 208. <http://dx.doi.org/10.1111/jfs.2007.27.issue-2>
- Analysis of the Association of Official Analytical Chemists. (2005). Official methods of analysis of the Association of Official Analytical Chemists (18th ed.). Gaithersburg, MD: Author.
- Analysis of the Association of Official Analytical Chemists. (2000). Official methods of analysis of the Association of Official Analytical Chemists (17th ed.). Gaithersburg, MD: Author.
- Larmond, E. (1991). Laboratory methods for sensory evaluation of foods (Series IV). Ottawa: Canadian Agricultural Publications.
- B.S. Anita, E.J. Akpam, P.A., Okon and I.U. Umoren, (2006) “Nutritive and antinutritive evaluation of Ipomoea batatas leaves pak,” *Journal of Nutrition*, 5(2): 166-168.
- Achinewhu SC (1986). The effect of fermentation on carbohydrate and fatty acid composition of African oil bean seeds. (*Pentaclethra macrophylla*). *Food Chem.*, 19: 105-116.
- Ndife, J., Obiegbuna, J.E., Akubor, P.I. and Ishiwu, C. N. (2013) Effect of substituting sugar with date palm pulp meal on the physicochemical, organoleptic and storage properties of bread. *Afr. J. Food Sci.* 7, 113–119
- Apena A, Opeolu SO, Bamidele FA, and Shittu A (2015). Nutrient changes during fermentation of some selected cereals. *Sky J Biochem. Res.* 4(2): 010 – 012.
- Makokha AO, Oniang’o RK, Njoroge SM and Kamar OK (2002). Effect of traditional fermentation and malting on phytic acid and mineral availability from sorghum (*Sorghum bicolor*) and finger millet (*Eleusine coracana*) grain varieties grown in Kenya. *Food and Nutrition Bulletin*, 23(3 Suppl), 241-245
- Slavin JL & Marlett JA (1997). American Dietetic Association: Health implication of dietary fiber. *J. Am. Diet. Assoc.* 97: 1157-1159
- Apena A, Opeolu SO, Bamidele FA, and Shittu A (2015). Nutrient changes during fermentation of some selected cereals. *Sky J Biochem. Res.* 4(2): 010 – 012.
- Odunfa SA and Adeyele S (1985). Microbiological changes during the traditional production of ogi-baba, a West African fermented sorghum gruel. *Journal of Cereal Science*, 3(2), 173-180.
- R.M. Costa, F.A.R. Oliveira, Modelling the kinetics of water loss during potato frying with a compartmental dynamic model, *Journal of Food Engineering*, Volume 41, 1999, Pages 177-185, ISSN 0260-8774, [https://doi.org/10.1016/S0260-8774\(99\)00095-3](https://doi.org/10.1016/S0260-8774(99)00095-3).
- Elkhalifa, A.E.O., and Bernhardt, R. (2013) Some physicochemical properties of flour from germinated sorghum grain. *J Food Sci Technol* 50, 186–190. <https://doi.org/10.1007/s13197-011-0315-2>
- Aremu M, Olaofe O, and Akintayo E (2007). Functional properties of some Nigerian varieties of legume seed flours and flour concentration effect on foaming and gelation properties. *J. Food Technol.* 5:109–115

- Mariscal, M. and P. Bouchon, (2008) Comparison between atmospheric and vacuum frying of apple slices, *Food Chemistry*, Volume 107, Issue 4, Pages 1561-1569, ISSN 0308-8146, <https://doi.org/10.1016/j.foodchem.2007.09.031>.
- Hart, M.R., Graham, R.P., Gee, M. and Morgan, A.I., JR. (1970), Bread from sorghum and barley flours. *Journal of Food Science*, 35: 661-665. <https://doi.org/10.1111/j.1365-2621.1970.tb04837.x>
- Taur, A. T., Pawar, V. D., & Ingle, V. M. (1984). Nutritional improvement of grain sorghum by germination. *The Indian Journal of Nutrition and Dietetics*, 21, 168– 173.
- Aly, M. and Seleem, H. (2015) Gluten-Free Flat Bread and Biscuits Production by Cassava, Extruded Soy Protein and Pumpkin Powder. *Food and Nutrition Sciences*, 6, 660-674. doi: 10.4236/fns.2015.6706
- Rosenthal, A. J. (2010). Texture profile analysis – How important are the parameters? *Journal of Texture Studies*, 41, 672–684.
- Szczesniak, A. S. (2002). Texture is a sensory property. *Food quality and preference*, 13(4), 215-225.
- Pilar Buera, M., Osella, C.A., Sánchez, H.D., Carrara, C.R., and de la Torre, M.A. (2005), Water Redistribution and Structural Changes of Starch During Storage of a Gluten-free Bread. *Starch/Stärke*, 57: 208-216. <https://doi.org/10.1002/star.200400330>
- Karaoğlu, M.M. and Kotancilar, H.G. (2009), Quality and textural behaviour of par-baked and rebaked cake during prolonged storage. *International Journal of Food Science & Technology*, 44: 93-99. <https://doi.org/10.1111/j.1365-2621.2007.01650.x>