

INTRODUCTION

Diabetes mellitus and thyroid diseases are two common endocrine disorder that affect multiple organs in the body and mutually affect each other (Wang *et al.*, 2018; Hage *et al.*, 2011). Thyroid hormones regulates carbohydrate metabolism and pancreatic function while diabetes mellitus affect TSH production, impairs the responds of TSH to TRH, reduced T3 level, and impairs the conversion of T4 to T3 (Hage *et al.*, 2011). On the other hand, Hyperthyroidism increases glucose production, absorption, and utilization thereby leading to hyperinsulinemia, abnormal glucose tolerance, and peripheral insulin resistance (Lambadiari *et al.*, 2011).

In developing countries, natural products especially from plants origin has been of great sources of relief for disease. One of such plant is *Adansonia digitata* with “Baobab” as English name. It belongs to the Malvaceae family. The different parts of the *A. digitata* provide food, shelter, clothing, medicine, and sources of raw materials for industrial use (Yakubu *et al.*, 2020; Venter and Venter, 1996). The antioxidant and anti-inflammatory properties of the tree makes it relevant in the treatment of different types of ailments (Kamatou *et al.*, 2011; De-Caluwe *et al.*, 2010). Recent phytochemical analysis of its leaves revealed that they contain a rich repertoire of reducing sugars, flavonoids, terpenoids, saponins, tannins, alkaloids, anthraquinones, steroids, resins, phenols, and cardiac-active glycosides (Abiona *et al.*, 2015; Finbarrs-Bello *et al.*, 2020). Traditionally in some parts of Africa, the leave is used to treat fatigue, dysentery, diseases of the urinary tract, ophthalmia and otitis (Abiona *et al.*, 2015). Moreover, it is used as a tonic, and for insect bites Guinea worm and internal pains, (Sidibe and Williams, 2002). Baobab leaves are potential source of protein used to complement amino acid to improve the overall protein quality in local diet (Nordeide *et al.*, 1996). The leaves are sources of minerals, iron and calcium (Barminas *et al.*, 1998; Boukari *et al.*, 2001). This study was aimed at determining the effects of aqueous extract of *Adansonia digitata* on the thyroid gland in alloxan-induced diabetic Wistar rats.

MATERIALS AND METHODS

Collection, Authentication and Preparation of plant materials

Fresh leaves of *A. digitata* were procured from a local dealer in Kaduna state. It was identified by a botanist in the Department of Plant Sciences and Biotechnology, University of Nigeria, Nsukka. The extraction method described by Bello and Nwoso (2015) was used. 500g of the powdered plant material was soaked in 1500ml of distilled water for 72 hours. Separation of the plant extract was carried out using Whatman NO. 4 filter paper. The resulting extract was then concentrated using rotatory evaporator, and stored at 4°C before use.

Experimental Animals and Ethical Approval

A total of 28 adult male Wistar rats were purchased from the animal house of the University of Nigeria, Enugu Campus. The animals were bred in the animal facility of Enugu State University of Science and Technology College of Medicine, Enugu. The animals were housed in netted iron cages and were provided easy access to food (Grower’s mesh, New market Enugu) and water *ad libitum*; and were maintained under standard laboratory conditions (temperature 20°C to 24°C, with relative humidity of 60-70% under 12 hours light and day cycles) and allowed to acclimatize for two weeks prior to the experiment. At

the end of acclimatization, the animals were randomly divided into seven (7) groups of four (4) animals each. They were labeled as:

Sham control - 0.1ml/kg/bw normal saline/14days

Diabetes control - Alloxan (150mg/kg)/2days

Diabetes + Low dose -150mg/kg/bw Alloxan +400mg/kg/bw *Adansonia digitata* extract

Diabetes + High dose -150mg/kg/bw Alloxan +600mg/kg/bw *Adansonia digitata* extract

Diabetes + Metformin -150mg/kg/bw Alloxan + 150mg/kg/bw Metformin

Low dose - 400mg/kg/bw *Adansonia digitata* extract only

High dose- 600mg/kg/bw *Adansonia digitata* extract only

The body weights of the animals were obtained before and after acclimatization, and at weekly intervals during the experiment. The protocol for conducting the in-vivo study in Wistar rats was approved by the Institutional Animal Ethical Committee (IAEC), Enugu State University of Science and Technology.

Induction of Diabetes using alloxan

Stock solution of alloxan was prepared by dissolving alloxan monohydrate (Sigma-Aldrich, USA) (0.9g) in 6ml of distilled water, and diabetes was induced by single intraperitoneal injection of alloxan monohydrate (150mg/kg). The volume of the solution containing 150mg/kg given to each rat was determined by its weight. After a period of two days, the rats with blood glucose level greater than 200mg/dl was considered diabetic and used for the research work. The method of Mohammad and Hauwa'u (2013) was adopted in the study with slight modification.

Hormonal assay

At the end of the experiment, blood samples were collected via cardiac puncture from all rats. Serum was then separated and stored at -20°C until the hormonal assay. Serum levels of thyroid hormones thyroxine and triiodothyronine concentrations were determined by ELISA (Enzyme-linked Immune-sorbent Assay) by Micallef *et al.* (1995)

Histopathological Investigation

Twenty four hours after the last treatment, the animals were sacrificed under anesthesia (ketamine hydrochloride). Sections of the thyroid were collected for histopathological examination. The samples were fixed in 10% phosphate buffered formalin for a minimum of 48 hours. The tissues were subsequently trimmed, dehydrated in 4 grades of alcohol (70%, 80%, 90% and absolute alcohol), cleared in 3 grades of xylene and embedded in molten wax. On solidifying, the blocks were sectioned, 5µm thick with a rotary microtome, floated in water bath and incubated at 60°C for 30 minutes. The 5µm thick sectioned tissues were subsequently cleared in 3 grades of xylene and rehydrated in 3 grades of alcohol (90%, 80% and 70%). The sections were then stained with Hematoxylin for 15 minutes. Blueing was done with ammonium chloride. Differentiation was done with 1% acid alcohol before counterstaining with Eosin. Permanent mounts were made on degreased glass slides using a mountant; DPX.

The prepared slides were examined with a Motic™ compound light microscope (Motic BA410E Elite Research Compound Microscope; Motic Asia, Hong Kong) using x4, x10

and x40 objective lenses. The photomicrographs were taken using a Motic™ 5.0 megapixels microscope camera at x160 and x400 magnification

Statistical Analysis

Data obtained was expressed as the mean standard deviation. They were analyzed using Statistical Package for Social Sciences (SPSS version 21; IBM SPSS, Chicago, Illinois, USA) software package. Values were analyzed as Mean± SD using one way ANOVA with Tukey post hoc test. The level of significance was set to $p < 0.05$.

RESULTS

Adansonia digitata increased triiodothyronine concentration

GROUPS	TRIODOOTHYRONINE (T3)	THYROXINE (T4)
Sham control	1.02±0.16	4.43±2.03
Diabetic control	1.02±0.16	2.42±1.32
Diabetic + Low dose	1.07±0.16	2.57±1.04
Diabetic + High dose	1.09±0.17	1.99±1.56
Diabetic + Metformin	0.57±0.22	0.87±0.27*
Low dose extract only	1.44±0.45 ^e	2.52±1.12
High dose extract on	1.47±0.43 ^e	2.60±1.11

KEYS: *P <0.05 compared with the control group 1; ^eP <0.05 compared with diabetic + metformin group.

The effect of *A. digitata* aqueous leaf extract on the serum concentrations of the thyroid hormones; triiodothyronine (T3) and thyroxine (T4) were analyzed. Significant increase was seen in triiodothyronine concentrations in low and high dose groups when compared to the diabetic + metformin group. Conversely, there was a significant decrease in thyroxine concentration in diabetic + metformin group when compared with sham control group.

Adansonia digitata reversed histological damage to the thyroid gland caused by diabetes

The effects of *Adansonia digitata* aqueous leaf extract on the histology of the thyroid gland was investigated. The thyroid gland of sham control, low- and high- dose animals showed normal histoarchitecture with parafollicular cells (arrow) and thyroid follicular cells observed. The parafollicular cells are mostly located within the basal aspects of the follicular cells and the basement membrane of the follicles. The follicles (F) are lined by cuboidal to columnar lining cells with their polarity directed towards the lumen of the follicles. The thyroid gland of the diabetic control animals showed vacuolar degeneration and random individual necrosis of the thyroid follicular cells and the parafollicular cells. The affected cells appear swollen with numerous minute clear intracytoplasmic vacuoles while the *A. digitata* and metformin treated groups recuperated and normal histo-

morphology of thyroid gland cells were observed. Follicles (F); Cells (White arrow). (H&E. x400)




Figure 1: Section of thyroid gland (A) Sham control - 0.1ml/kg normal saline/14days

(B) Diabetes control (Alloxan, 150mg/kg). (C) Diabetes + Low dose -150mg/kg/bw Alloxan +400mg/kg/bw *Adansonia digitata* extract (D) Diabetes + High dose -150mg/kg/bw Alloxan +600mg/kg/bw *Adansonia digitata* extract/bw (E) Diabetes + Metformin -150mg/kg/bw Alloxan + 150mg/kg/bw Metformin (F) Low dose -400mg/kg/bw *Adansonia digitata* extract only. (G) High dose- 600mg/kg/bw *Adansonia digitata* extract only.

DISCUSSION

In this study, we used the leaf extract of *Adansonia digitata* on alloxan-induced diabetic rats to investigate the relationship between *A. digitata* leaves extract and thyroid gland of diabetes mellitus Wistar rats with an aim to establish its usefulness in the treatment of diabetes with co-morbid thyroid dysfunction. Sathish and Mohan (2003), revealed that diabetes may induce a “low T3 state”. In contrast to this findings, differences were not found in the serum levels of triiodothyronine (T3) of both sham control and diabetic control groups. However, animals that received *A. digitata* had increase T3 even though the increase was not significant when compared to the sham control and diabetic control. Significant differences was observed in groups that receive *A. digitata* without diabetes and diabetic + metformin group. This implies that *A. digitata* had potency to increase serum triiodothyronine (T3) levels and also had the capacity to restore the hormone back to its normal levels in diabetic conditions. More proof of this fact could be seen in the histology of the thyroid gland where the medium and high dose of the aqueous leaf extract of *Adansonia digitata* were sufficient to reverse the thyroid toxicity brought about by Alloxan administration.

Although diabetes and *A. digitata* led to reduction of thyroxine (T4) level, significant differences were not found in serum levels of T4 in all groups except in diabetic + metformin group when compared with the sham control suggesting that metformin has the ability to reduce serum T4 level. The reduction in T4 level of animals treated with metformin contrast previous works by Hu *et al.* (2017) where metformin led to significant increase in T4 level of animals and Anil *et al.*(2016) where changes was not observed among metformin treated and non-treated animals.

CONCLUSION

The present study indicates that treatment of alloxan-induced diabetes rats with *Adansonia digitata* extracts increased thyroid hormones concentrations, and restored the diabetic thyroid gland to its normal state.

CONFLICT OF INTEREST

Authors declared they have no competing interests

AUTHORS' CONTRIBUTIONS

Elizabeth Finbarrs-Bello designed and wrote the protocol for the study, OPO and NMC wrote the first draft of the manuscript. Authors OPO and EFB managed the analyses of

the study. Authors NMC and EFB managed the literature searches. All authors read and approved the final manuscript.

FUNDING: Self Funded

REFERENCES

Wang C, Dai X, Zhang D, Liu Z, Huang Q (2018). Jinlida Granules Improve Dysfunction of Hypothalamic-Pituitary-Thyroid Axis in Diabetic Rats Induced by STZ. *BioMed Research International*

Hage M, Zantout MS, Azar ST (2011). Thyroid disorders and diabetes mellitus. *Journal of Thyroid Research*

Lambadiari V, Mitrou P, Maratou E et al (2011). Thyroid hormones are positively associated with insulin resistance early in the development of type 2 diabetes. *Endocrine Journal*; 39(1): 28-32.

Venter F and Venter JA (1996). Baobab. In: Making the Most of Indigenous Trees, Venter F and JA Venter (Eds.), Briza Publications, Pretoria, South Africa. 26-27.

Yakubu OE, Imo C, Shaibu C, Akighir J, Ameh DS. (2020). Effects of Ethanolic Leaf and Stem-bark Extracts of *Adansonia digitata* in Alloxan-induced Diabetic Wistar Rats. *J. Pharmacol. Toxicol.* 15(1): 1-7.

Kamatou GPP, Vermaak I, Viljoen AM (2011). An updated review of *Adansonia digitata*: A commercially important African tree. *South African Journal of Botany*, 77(4): 908-919.

De Caluwe E, Halamova K, Damme PV (2010). Baobab (*Adansonia digitata* L.): a review of traditional uses, phytochemistry and pharmacology. *Atuo Focus*. 23:11-51.

Abiona DL, Adedapo Z, Suleiman MK (2015). Proximate analysis, phytochemical screening and antimicrobial activity of baobab (*Adansonia digitata*) leaves. *IOSR Journal of Applied Chemistry*. 8: 60-65

Finbarrs-Bello E, Neboh EE , Nnaji KS , Nwifor GO (2020) Effects of Aqueous leaf extract of *Adansonia Digitata* (baobab) on alloxan-induced diabetes mellitus on endocrine pancreas of adult male Wistar rats. *Journal of experimental research*. 8(3):23-34.

Sidibe M, William JT (2002). Baobab *Adansonia digitata* L. in: Hughes A, Smith R.W, editors fruits for the future 4 Southampton uk. *International centre for underutilized crops*. P: 96.

Nordeide M, Hatloy A, Folling M, Oshaug E (1996). Nutrient composition and Nutritional importance of green leaves and wild food sources in an agricultural district, Koutiala, Southern Mali. *International Journal of Food science and Nutrition*. 47: 455-468.

Barminas JT, Charles M, Emmanuel D(1998). Mineral composition of non-conventional leafy vegetables. *Plant Foods for Human Nutrition*. 53: 29-36.

Boukari I, Shier NW, Fernandez RXE, Frisch J, Watkins BA, Pawloski L, Fly AD (2001). Mineral Analysis of Selected Western African Foods. *Journal of Food Composition and Analysis*. 14:37-42.

Bello OK and Nwoso SO (2015). Radioactive potentials of methanolic/aqueous extract of *Adansonia digitata* and *Cochorous olitorious* leaf plant on gamma irradiated male Wistar rats. *Journal of Natural Sciences Research*; 5(11): 81-93

Sathish R and Mohan V (2003). Diabetes and thyroid diseases - A review. *International Journal For Diabetes In Developing Countries*. 23: 120-123.

Hu X, Liu Y, Wang C, Hou L, Zheng X, Xu Y, Ding L, Pang S (2017). Metformin affects thyroid function in male rats. *Oncotarget*. 8(64): 107589-107595

Anil C, Kut A, Atesagaoglu B, Nar A, Tuntuncu NB, Gursoy A (2016). Metformin Decreases Thyroid Volume and Nodule Size in Subjects with Insulin Resistance: A Preliminary Study. *Med Princ Pract*. 25:233-236.