

CHEMICAL ANALYSIS AND *IN-VITRO* ANTIOXIDANT POTENTIAL OF *DATURA STRAMONIUM* FRUIT-OIL GROWN IN OGBOMOSO SOUTHWEST NIGERIA

Fatoki JO^{*1}, Badmus JA², Adekomi DA³, Atere TG¹,
Ajilore BS¹, Kehinde SA⁴, Faokunla O⁵

¹Department of Medical Biochemistry, Faculty of Basic Medical Sciences, College of Health Sciences, Osun State University, Osogbo, Nigeria.

²Department of Biochemistry, Faculty of Basic Medical Sciences, College of Health Sciences, Ladoke Akintola University of Technology, Ogbomosho, Nigeria.

³Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, Osun State University, Osogbo, Nigeria.

⁴Department of Chemical Science, Faculty of Natural Sciences, Ajayi Crowther University, Oyo, Nigeria.

⁵Department of Biochemistry, Federal University of Lokoja, Lokoja, Nigeria.

*Author for Correspondence: john.fatoki@uniosun.edu.ng

ABSTRACT

The present study analyzed the chemical components and the antioxidant potentials of and antimicrobial activities of the oil isolated from *Datura stramonium* fruits that were grown in Ogbomosho, Southwest, Nigeria. Fresh fruits of *Datura stramonium* were collected in the premises of LAUTECH Teaching Hospital, Ogbomosho, South West Nigeria around February 2018. The fruit oil was thereafter extracted with 70% methanol and concentrated with a rotary evaporator. The iodine, saponification, acid, peroxide values were determined titrimetrically. An *in-vitro* 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity, total phenol content, reductive potential, ferric reducing antioxidant assay, and hydroxyl radical scavenging activity was used to determine the antioxidants activity of the oil spectrophotometrically. Gas chromatography-mass spectrophotometry (GC-MS) and high-performance liquid chromatography (HPLC) were used to identify the chemical constituents of the oil. The iodine, saponification, acid, and peroxide values of the oil were 86.15 ± 1.01 mg/g oil, 253.85 ± 0.98 MgKOH/g, 5.27 ± 0.032 mgKOH/g, and 2.80 ± 0.79 milli – equivalentO₂/kg respectively. A concentration-dependent increase in *in-vitro* antioxidant activities characterized the data from all the antioxidant parameters tested. While results obtained from GC-MS analysis reveals that the *Datura stramonium* fruit oil contained phenolic compounds, alkene compounds, aliphatic compounds, amide compounds, and fatty acid, HPLC identified the presence of catechin, rutin, and kaempferol, Oil from *Datura stramonium* fruit has potent antioxidant potentials and can therefore be looked into for possible utilization.

Keywords: antioxidants, chromatography, *Datura stramonium*, free radicals, fruit oil.

INTRODUCTION

Datura stramonium is also known as Jimson weed, Locoweed, Angel's trumpet, Devil's trumpet. Its common name is Africa Thorn apple and Gegemu in Yoruba. It is a hallucinogenic plant found in the urban and rural areas, in cornfields, and along road sides (Bouziri et al. 2011). *Datura* genus is widely distributed across the tropical and warm temperate regions of the world. *Datura stramonium* is native to North America, Europe, Asia, and Africa (Parashuram, 2011). It is germinated through fruits. The germination period of the fruit is usually 3 to 6 weeks. When properly stored, the fruits will remain viable even after several years. *D.*

stramonium is an annual flowering plant. It herbaceous stem, with branched, and glabrous or only slightly hairy. When cultivated, a fully grown *Datura stramonium* plant can be up to about 1 meter tall (Norton, 2010). The stems of *Datura stramonium* are highly branched and are spreading; the stems are also stout, leafy, smooth, and erect, with pale yellowish-green in appearance. Leaves are simple dentate, slightly hairy, oval glabrous, bid, and about 4 to 6 inches long. *Datura stramonium* bears funnel-shaped, white, or purple-colored flowers. The average length of the flower is about 3 inches. Fruits are large and full of thorns (Norton, 2010).

About ten species of *Datura* are known, out

of which *Datura anoxia* and *Datura stramonium* are mostly researched into with potent phytochemical of pharmacologic importance. It had been reported that every part of the *Datura stramonium* plant, such as the leaves, stems, flowers, fruits, and even the root are all capable of producing secondary metabolites that are of medical importance (Elgohary et al. 2018). Researchers are lately showing more and more interest in the effective biological potentials of oils, especially those of plant origin. These biological activities include but are not limited to their antioxidant capacities (Elgohary et al. 2018). Furthermore, increasing data from previous studies have suggested the usage of oils in the food and drug industries as a source of natural antioxidants due to their potential antioxidant capacities and extremely low toxic potentials. The antioxidants and other biological activities of many of these oils from plant origin can be attributed to the presence of the plant's secondary metabolites such as phenol, flavonoids, and others (Asari et al. 2019).

Consequently, the present study aimed to assess the quality and the antioxidant potentials of *Datura stramonium* fruit oil.

MATERIAL AND METHODS

Chemicals

Chemicals and reagents used for this study were obtained from British Drug House (BDH) Chemicals Limited Poole England and Sigma-Aldrich, Missouri, U.S.A. all reagents were of analytical grade.

Plant material

Fresh fruits of *Datura stramonium* were collected in the premises of LAUTECH Teaching Hospital, Ogbomoso, South West Nigeria around February 2018, and were authenticated by Prof. A.T. J. Ogunkunle, of the Department of Pure and Applied Biology, Faculty of Pure and Applied Sciences Ladoke Akintola University of Technology, Ogbomoso, Nigeria, with voucher number PAB/19/176.

Collection and extraction procedure

Datura stramonium fruits were washed with clean water, cut into small pieces to facilitate the drying process, and were allowed to air-dry under room temperature after which the dried fruit was

ground to a very fine powder with the aid of an electric blender, and extracted as described by Kumar et al. (2010).

The powdered fruits of *Datura stramonium* (2 kg) were exhaustively extracted with 6 liters of 70% methanol. The methanolic filtrate was then concentrated with the aid of a rotary evaporator at 60°C to obtain a dark-brown oily substance.

Determination of free fatty acid value

Free fatty acid value of the fruit oil was determined according to the official method of American Oils Chemists' Society (AOCS, 2004). 10 g of the sample was weighed into a conical flask containing 50 ml ethanol that had been previously neutralized by the addition of 2 ml phenolphthalein solution. The mixture was thoroughly vortexed and placed in a water bath at 40°C. The mixture under this condition is gently titrated with 0.25 N Sodium hydroxide (NaOH) with vigorous shaking until a permanent faint pink color that persists for at least 1 minute is developed to obtain the titer value. The free fatty acid value of the sample was calculated from the titer value using the following formula:

$$\text{FFA} = \frac{28.2 \times N \times V}{W}$$

Where:

N = normality

V = volume NAOH used (titre value)

W = weight of the sample

Determination of saponification value

Saponification value of the fruit oil was determined according to the official method of American Oils Chemists' Society (AOCS, 1990). 25 mL of potassium hydroxide/ethanol was added to 2 g of the test sample in a conical flask with an attached reflux condenser. The mixture was heated continuously in a boiling water bath for 30 minutes to allow all the fat to saponify completely. 1 mL of phenolphthalein/ethanol was added to the mixture immediately after boiling and the excess of alkali was titrated with 0.5 N hydrochloric acid and the titer value was noted. A blank test containing the same quantities of reagents but without the test sample was equally carried out

and the titer value was also noted.

The saponification value of the sample was calculated from the titer values using the following formula:

$$\text{Saponification value} = \frac{(V_2 - V_1) \times 0.028805 \times 1000}{\text{weight(g) of the test sample}}$$

Where:

V_1 = titer value of the sample

V_2 = titer value of the blank

Determination of iodine value

The iodine value of the sample was done using the method of Wijs (Pearson, 1970). The Wijs' reagent was prepared by dissolving 8.0 g iodine trichloride (ICl_3) in 200 mL glacial acetic acid and 9.0 g iodine in 300 mL CCl_4 . The two solutions were mixed and diluted to 1 L with glacial acetic acid. 0.3 g of the test sample was weighed and dissolved in carbon tetrachloride (CCl_4) and 25 mL of Wijs' reagent was added. The mixtures were thoroughly mixed and allowed to stand for 30 minutes in the dark. 20 ml of 10% potassium iodide solution and 100 mL distilled water were added. The iodine liberated was titrated with 0.1 M sodium thiosulphate solution using starch as the indicator just before the endpoint. A blank test containing the same quantities of reagents but without the test sample was equally carried out. The iodine value of the sample was calculated from the titer values using the following formula:

$$\text{Iodine value (mg I}_2\text{/100 g sample)} = \frac{(V_1 - V_2) \times 1.269}{W}$$

Where:

V_1 = volume (cm^3) of thiosulphate used to titrate blank

V_2 = volume (cm^3) of thiosulphate used to titrate test sample

W = weight (g) of the test sample.

Determination of peroxide value

Peroxide value of the fruit oil was determined according to the official method of American Oils Chemists' Society (AOCS, 2005). 10 ml of acetic acid chloroform solution (ratio 4/6) was added to 0.3g of the test sample in a 250 ml flask with a stopper shaking vigorously to allow the test sample to dissolve after which 1 ml of potassium iodide was added and the entire mixture was

allowed to stand in the dark for 5 minutes at room temperature. 20 ml of distilled water was thereafter added with continuous shaking. The liberated iodine was titrated with 0.01 N sodium thiosulphate solution and once a light yellow coloration was observed 1 ml of 1.5% starch solution was added as an indicator and titrated till the mixture turned colorless. A blank test containing the same quantities of reagents but without the test sample was equally carried out. The peroxide value of the sample was calculated from the titer values using the following formula:

$$\text{Peroxide value (meq/kg)} = \frac{(V_1 - V_2) \times C \times 1000 \times N}{W}$$

Where:

V_1 = titer value of the sample

V_2 = titer value of the blank

C = molar concentration of sodium thiosulphate

N = normality of sodium thiosulphate

W = weight of the test sample.

1,1-Diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging activity

DPPH free radical scavenging activity of the oil was assayed using the DPPH photometric method of Mensor et al. (2001). When an antioxidant compound that is capable of donating hydrogen reacts with DPPH, the DPPH is reduced with a concomitant change of color from deep violet to golden/light yellow can be monitored at 518 nm. Briefly, 1 mL of 0.3 mM of DPPH solution was added to 1 ml of the oil sample, the resultant mixture was incubated at room temperature in the dark for 30 min. The absorbance values were read at a wavelength of 518 nm. The percentage antioxidant activity was then calculated from the absorbance, using the formula below:

$$\text{DPPH scavenging effect (\%)} = \left(\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \right) \times 100$$

Total phenol content

This was estimated as described by McDonald et al. (2001). The assay is based on the reduction of Folin-Ciocalteu reagent by the phenolic compounds. The reduced Folin-Ciocalteu reagent is blue and the absorbance can be measured at the wavelength of 500-750 nm. Briefly, 1 ml of the oil sample was added to 0.2

mL each of Folin-Ciocalteu reagent and 2 mL of distilled water. One mL of 15% Na_2CO_3 was mixed with the solution. The mixture was incubated for 30 minutes at 40°C and the absorbance was thereafter read at 760 nm. The total phenol content of the sample was expressed as $\mu\text{g}/\text{mg}$ of gallic acid equivalent (GAE).

Reductive potential

The method of Oyaizu (1986) was employed for the assessment of the reductive potential of the oil sample. 150 μL of the oil sample in 1 mL of distilled water was mixed with 2.5 mL phosphate buffer (0.2 M, pH 6.6) and 1% potassium ferricyanide. The mixture was incubated at 50°C for 20 min. 2.5 mL of 10% trichloroacetic acid (TCA) was added and the solution was centrifuged at 1000 X g for 10 min. 2.5 mL of the upper layer of solution was thereafter mixed with 2.5 mL of distilled water and 0.5 mL of 1% FeCl_3 . The absorbance was read at 700 nm.

Ferric reducing antioxidant power assay (FRAP)

FRAP assay was done according to the method described by Tsai et al. (2011), with slight modification. FRAP uses antioxidants as reductants in a redox-linked colorimetric method to test the total antioxidant power directly. A reagent containing 0.83 mmol L⁻¹ TPTZ and 1.67 mmol L⁻¹ ferric chlorides at 0.1 mol L⁻¹ acetate buffer (pH 3.6). The sample was mixed with 0.9 ml of reagent and incubated at 25°C for 10 min. Formation of a blue-colored ferrous tripyridyltriazine (Fe^{2+} -TPTZ) from a yellow ferric tripyridyltriazine (Fe^{3+} -TPTZ) was observed at an absorbance of 593 nm. Values were expressed in $\mu\text{mole Fe}^{2+}/\text{mg}$ dry weight of the test sample. Gallic acid was used as the standard reference.

Hydroxyl radical (OH) scavenging activities

All solutions were freshly prepared before the assay. One (1) mL of the reaction mixture contained 100 μL of 2.8 mM 2-deoxyribose (dissolved in phosphate buffer (10 mM), pH 7.4), 500 μL solution of various concentrations of the extract (25, 50, 75 and 100%), 200 μL of 200 μM FeCl_3 and 1.04 μM EDTA (1:1 v/v), 100 μL of H_2O_2 (1.0 mM) and 100 μL of ascorbic acid (1.0

mM). After an incubation period of 1 hour at 37°C , the extent of deoxyribose degradation was measured by TBA reaction (Halliwell et al. 1987). The % inhibition of hydroxyl radical was calculated and calculation of results as described above.

Gas Chromatography-Mass Spectroscopy

The GC-MS analysis was carried out according to the method described by Ajayi et al. (2011).

High-Performance Liquid Chromatography (HPLC)

HPLC was carried out using an apparatus consisting of a pump (Model LC10AT and FCV-10al) and DGU-14A degasser (Shimadzu, Tokyo, Japan), equipped with a UV-VIS detector model L-4250 (Merch Hitachi, LTD Tokyo, Japan). Data acquisition was performed by an interface Model Data Apex CSW32 (Prague, Czech Republic) with a PC. A reversed-phase Supelcosil column LC18 (15 cm x 4.6mm, ID 3 mm particle size), and a Supelguard cartridge column (Supelco, Bellafonte, PA) were used for the chromatographic analysis. The absorbance of the eluted compounds was monitored at 242 nm. The mobile phase was acetonitrile-water 85 –15 (v/v) at a flow rate of 1 mL/min. The peaks were identified by their retention times in comparison to the external standards.

Statistical Analysis

All the experimental results were the mean \pm SEM of three independent measurements, statistically analyzed using one-way analysis of variance (ANOVA) followed by Dunnett's *t*-test using Graphpad® prism version 5.02 statistical software.

RESULTS

Table 1 showed the iodine value, saponification value, free fatty acid value, and the peroxide value of *Datura stramonium* fruit oil. As presented value for each of the parameter determined were 86.15 ± 1.01 mg/g oil, 253.85 ± 0.98 MgKOH/g, 5.27 ± 0.032 mgKOH/g, and 2.80 ± 0.79 milli – equivalent O_2/kg respectively.

Table 1. Chemical parameter of *Daturastramonium* fruit oil

S/N	Parameters	Results
1	Iodine Value	86.15 ± 1,01 mg/g oil
2	Saponification Value	253.85 ± 0.98 MgKOH/g
3	Free Fatty Acid Value	5.27 ± 0.032 mgKOH/g
4	Peroxide Value	2.80 ± 0.79 milli- equivalentO ₂ /kg

Values are expressed as the mean of triplicate measurements ± standard error of the mean (SEM)

Table 2 depicts the mean value of percentage DPPH scavenging activities of *Datura stramonium* fruit oil. The fruit oil caused a concentration-dependent increase in the percentage of DPPH radical scavenging activities. For instance, the lowest concentration of 25% led to 28.37 ± 0.021% DPPH radical scavenging activity, while the highest concentration of 100% induced 82.57 ± 0.19% DPPH radical scavenging activities.

Table 2. Percentage DPPH scavenging activities of *Datura stramonium* fruit oil

Concentration	% DPPH scavenging activities
25%	28.37 ± 0.021
50%	50.48 ± 0.017
75%	68.84 ± 0.09
100%	82.57 ± 0.19

Values are expressed as the mean of triplicate measurements ± standard error of the mean (SEM)

The mean value of total phenol contents of *Datura stramonium* fruit oil using garlic acid as the standard was presented in Table 3. As showed, the fruit oil contained an appreciable quantity of phenolic compounds. The data indicated that the contents of the total phenol in the fruit oil increased as the concentration of *Datura stramonium* fruit oil tested increased. That is the lowest concentration of 25% contained the lowest amount of phenolic compounds (0.170 ± 0.003 µg/mg GAE), while the highest concentration of 100% contained the highest amount of total phenols (0.238 ± 0.009 µg/mg GAE).

Table 3. Total phenol contents of *Datura stramonium* fruit oil

Concentration	Total phenol content (µg/mg GAE)
25%	0.170 ± 0.003
50%	0.197 ± 0.001
75%	0.210 ± 0.017
100%	0.238 ± 0.009

Values are expressed as the mean of triplicate measurements ± standard error of the mean (SEM)

The data obtained from the assessment of reductive potentials of *Datura stramonium* fruit oil are presented in Table 4 expressed as mean value \pm SEM. The data presented showed that the *Datura stramonium* fruit oil possessed reductive

potentials when tested *in vitro*. As shown, 25, 50, 75, and 100% *Datura stramonium* fruit oil when tested *in vitro* potential power in the order $0.129 \pm 0.03 \mu\text{g/ml}$, $0.150 \pm 0.02 \mu\text{g/ml}$, $.0170 \pm 0.05 \mu\text{g/ml}$ and $1.110 \pm 0.008 \mu\text{g/ml}$ respectively.

Table 4. The reductive potential of *Datura stramonium* fruit oil

Concentration	Reductive potential ($\mu\text{g/ml}$)
25%	0.129 ± 0.03
50%	0.150 ± 0.02
75%	$.0170 \pm 0.05$
100%	1.110 ± 0.008

Values are expressed as the mean of triplicate measurements \pm standard error of the mean (SEM)

The mean value of ferric reducing antioxidant power of *Datura stramonium* fruit oil using garlic acid as a reference antioxidant is depicted in Table 5. As presented, *Datura stramonium* fruit oil at 25% concentration showed 66.78 ± 0.18

mg/ml ferric reducing antioxidant power, while at 50, 75 and 100%, the antioxidant power increased to $81.32 \pm 0.79 \text{ mg/ml}$, $89.26 \pm 0.35 \text{ mg/ml}$ and $121.33 \pm 0.48 \text{ mg/ml}$ respectively.

Table 5. Ferric Reducing Antioxidant Power (FRAP) of *Datura stramonium* fruit oil

Concentration	Mean reducing antioxidant power (mg/ml)	Reference antioxidant
25%	66.78 ± 0.18	Gallic acid
50%	81.32 ± 0.79	Gallic acid
75%	89.26 ± 0.35	Gallic acid
100%	121.33 ± 0.48	Garlic acid

Values are expressed as the mean of triplicate measurements \pm standard error of the mean (SEM)

The percentage hydroxyl radical scavenging activity of various concentrations of *Datura stramonium* fruit oil was presented in Table 6. The fruit oil when tested *in vitro* showed 65.08 ± 0.0034 percentage hydroxyl radical scavenging activities at 25%, 76.07 ± 0.009 percentage

hydroxyl radical scavenging activities at 50% and 83.14 ± 0.006 , 98.78 ± 0.005 percentage hydroxyl radical scavenging activities at 75% and 100% respectively indicating a concentration-dependent increase hydroxyl radical scavenging activities of the *Datura stramonium* fruit oil.

Table 6. Percentage hydroxyl radical scavenging activity of *Datura stramonium* fruit oil

Concentration	% hydroxyl radical scavenging activity
25%	65.08 ± 0.0034
50%	76.07 ± 0.009
75%	83.14 ± 0.006
100%	98.78 ± 0.005

Values are expressed as the mean of triplicate measurements \pm standard error of the mean (SEM)

As presented in Table 7, GC-MS identified compound, amide compound, and fatty acids in phenolic compound, alkene compound, aliphatic *Datura stramonium* fruit oil.

Table 7. Chemical compounds identified by GC-MS in *Datura stramonium* fruit oil

PeakNumber	Retention time (min)	Area (%)	Group of a compound identified
16	9.861	1.18	Phenolic compound
17	10.104	40.89	Phenolic compound
18	10.208	13.28	Phenolic compound
22	10.756	9.08	Alkene compound
23	11.080	2.54	Alkene compound
27	11.842	12.89	Phenolic compound
31	12.414	1.13	Alkene compound
35	13.095	2.27	Aliphatic compound
52	17.063	1.19	Fatty acid
57	18.703	2.42	Amide compound





Figure 1. Gas chromatogram of *Datura stramonium* fruit oil

As showed, Table 8 indicated that three different compounds were identified in the *Datura stramonium* fruit oil with the aid of high-

performance liquid chromatography. The three compounds as identified by HPLC are: catechin, rutin, and kaempferol.

Table 8. Compounds identified by HPLC in *Datura stramonium* fruit oil

Retention time (min)	Amount (PPM)	Identified compound
1.935	402.73788	Catechin
2.502	241.10211	Rutin
9.503	2.38724	Kaempferol

**Figure 2. HPLC chromatogram of *Datura stramonium* fruit oil**

DISCUSSION

Previous researches have shown that consumption of oils has diverse effects on cellular lipid homeostasis and the development of chronic disease, and well-being. Since oil from various sources generally have different composition and none of the oil is suitable for all purposes. Researchers have therefore beamed their searchlight on new sources of edible oils especially from plant fruits which are known to be a good source of healthy oils. In this study, chemical characteristics, phenolic content, and antioxidant activity of *Datura stramonium* fruits oil were determined (Ramadan et al. 2006).

Meanwhile, despite the acclaimed health benefits of oil from plant origin, it has been suggested that consumption of some of these oils may not be suitable for consumption as their consumption by a human may lead to the onset of cardiovascular diseases (AlKhattaf et al. 2020).

In the present research, we did a preliminary assessment of the quality of *Datura stramonium* fruits oil by the determination of its iodine value, saponification value, acid value, and peroxide value.

Iodine value is the most often quoted value of oil. It is a measure of the degree of unsaturation of oil. It is defined as the amount of iodine (in g) added to 100g of oil. Since the oxidative stability and melting point are related to the degree of unsaturation, the iodine value, therefore, gives an estimation of these qualities. The greater the iodine value, the more the degree of unsaturation and the higher the susceptibility of the oil to oxidation. That is, oils with high **iodine values** contain more of the highly unsaturated fatty acids that are prone to fast degradation reactions such as autoxidation or polymerization (Dijkstra, 2015). The observed high iodine value in the present study (86.15 ± 1.01 mg/g oil), depicted

that *Datura stramonium* oil has a high concentration of unsaturated fatty acids and possibly a low degree of saturation. Meanwhile, previous studies have recommended a switch from saturated fats to unsaturated fats because of the high risk of a cardiovascular disorder that is often associated with high consumption of saturated fatty-acids. Meanwhile, highly unsaturated oils can, however, undertake oxidative degradation as a result of their double bonds, except sufficient antioxidant is added (Adetola et al. 2016).

The saponification value indicates the molecular weights of triacylglycerol in an oil sample. The saponification value has an inverse relationship with the average chain length or weight of the fatty acids in the oil. That is, a higher Saponification value depicts a high amount of fatty acids with a lower carbon chain. Therefore, the shorter the average carbon length (C_4 - C_{12}) the higher is the saponification value (Muhammad et al. 2011). The high value obtained for *Datura stramonium* oil in the present study (253.85 ± 0.98 MgKOH/g) depicts that *Datura stramonium* oil extract contains high amounts of short-chain fatty acids ($< C_{12}$). According to Codex Alimentarius (2005) and the Asian and Pacific Coconut Community (APCC) Standards, the standard permissibility level of saponification values ranges between 250 – 260 mg KOH/g oil and 248 – 268 mg KOH/g oil respectively. This means that the saponification value of *Datura stramonium* obtained in the present study fell within the Codex and APCC standard permissible level and thus meets up with the required standard. The high saponification value indicates that *Datura stramonium* oil has the potential to be used in the cosmetic industry, and may also be suitable for soap making (Akinhanmi and Atasie, 2008).

The fatty acid value is the number of milligrams of potassium hydroxide (KOH) that is required to neutralize the fatty acids in 1 gram of sample. It is a measure of the amount of free fatty acids in the oil. Normally, fatty acids are found in the triacylglycerol form, however, during extraction/processing the fatty acids may get hydrolyzed into free fatty acid and glycerol in a reaction catalyzed by lipase. The higher the value of fatty acid of oil, the higher the level of free fatty

acids which is an indication of decreased oil quality. According to the American Oil Chemist Society, the acceptable levels for all oil samples should be ≤ 0.6 mg KOH/g. The acid values obtained in this study for *Datura stramonium* were 5.27 mg KOH/g, which is above the permissible level of 0.6 mg KOH/g. The high acid value displayed by *Datura stramonium* oil depicted that most of the triacylglycerol contained in the oil had been hydrolyzed to free fatty acids which suggest reduced oil quality (Mozaffarian and Clarke, 2009).

The peroxide value of any oil determines its oxidative state. It is equally an important marker in determining the quality of the oil. Since peroxide is the early product of oxidation, the peroxide value of oil provides information on how fresh is the oil. Elevated peroxide value is an indication that lipid oxidation had taken place. A peroxide value that is more than 9 meq O_2 /kg is an indication of a bad oxidation state of the oil. As reported in this study, a peroxide value of 2.8 – equivalent O_2 /kg indicates that the *Datura stramonium* oil used in this study is in a good state and has not been oxidized which represents good extraction and maintenance conditions (Dijkstra, 2015). This means that *Datura stramonium* fruits oil can be stored for a long time without becoming rancid as a result of oxidation. Meanwhile, a peroxide value of between 20 and 40 mill-equivalent O_2 /Kg indicates that the oil is rancid and oxidatively damaged. Furthermore, the Peroxide value of *Datura stramonium* oil is even lower than that of pomegranate fruits oil (3.42 mill-equivalent O_2 /Kg), linseed oil (11.28 mill-equivalent O_2 /Kg), and sunflower oil (12.87 mill-equivalent O_2 /Kg) (Dijkstra, 2015).

Antioxidant generally refers to a compound that is capable of neutralizing free radicals typically by donating an electron to the free radicals to make the free radical to be more a stable compound. In other words, when an antioxidant compound donated an electron to the reactive radicals (free radicals), it will convert the radicals into a more stable and passive condition and thereby aborting the chain reaction of the free radical (Ganu et al. 2010). The ability of *Datura stramonium* to function as an antioxidant was in this study access through its

percentage DPPH scavenging activities, reductive potential, ferric reducing antioxidant power, percentage hydroxyl radical scavenging activity, and total phenol contents.

DPPH is a free radical that is stable at room temperature; it produces a purple coloration in methanol. In the presence of an antioxidant molecule, it is reduced and gives rise to a yellowish methanol solution. The ability of a molecule to donate a hydrogen atom to a free radical is one of the mechanisms involved in the evaluation of antioxidant activity (Badmus et al. 2011). As shown, the DPPH radical scavenging activities of *Datura stramonium* fruit oil (Table 2) showed that the *Datura stramonium* possesses high antioxidant activity. This activity increases with increasing concentration in a dose-dependent manner. This observed antioxidant potential may be due to the presence of phenolic compounds in the *Datura stramonium* oil as observed in this study.

The reducing power of a biological compound is related to its ability to transfer electrons, and it may therefore be used as an important marker of its likely antioxidant potential (Sanchez-Moreno, 2002). As presented in Table 4, the methanolic extract of *Datura stramonium* showed a significant reductive capability. This reducing capability increased with increasing concentration of the *Datura stramonium* fruit oil. This finding implies that extract of *Datura stramonium* possesses hydrogen ion (H^+) donating potential and as such has the potential to serve as an antioxidant. This finding is in total agreement with every other obtained in the present study.

The total antioxidant power of a plant extract can be assessed by Ferric Reducing Antioxidant Power assay (FRAP) (Szollosi and Varga, 2002). FRAP assay depends upon the ferric tripyridyltriazine (Fe(III)-TPTZ) complex to the ferrous tripyridyltriazine (Fe(II)-TPTZ) by a reductant at low pH. Fe (II)-TPTZ is a substance with an intensive blue coloration that can be monitored spectrophotometrically at 593 nm. Data from previous studies reveal that FRAP assay is highly sensitive for the determination of the total antioxidant potential of the various biological substances, such as plant extracts. In the FRAP method, the antioxidants in the

biological compounds serve as a reductant in a redox-linked colorimetric reaction (Szollosi and Varga, 2002). In this study, the antioxidant capacity of the *Datura stramonium* oil was measured as ferric reducing antioxidant power (FRAP). Data obtained revealed that oil extracted from the plant possesses great antioxidant capacity. This is not surprising as it is in total agreement with other data obtained in the present study.

The hydroxyl radical is one of the most dangerous free radicals produced in cells (Medda et al. 2021). It is the most reactive product of reactive oxygen species generated from successive single electron reductions of molecular oxygen (O_2) during metabolic processes in the cell and is primarily responsible for the cytotoxic effects that are observed in aerobic organisms. It is generally assumed that OH is generated in biological systems from H_2O_2 by the Fenton reaction (Medda et al. 2021). It is considered to be one of the rapid initiators of the lipid peroxidation process; they do this by abstracting hydrogen atoms from polyunsaturated fatty acid (PUFA), which causes the peroxidation of membrane lipids. Hydroxyl radicals also abstract hydrogen atoms from each of the carbon atoms of the sugar moiety of DNA leading to oxidative damage of DNA (Badmus et al. 2011), and these effects have been implicated in mutagenesis, carcinogenesis, and aging (Halliwell and Gutteridge, 1999).

In this study, methanolic extract of *Datura stramonium* fruit was found to cause concentration-dependent hydroxyl radical scavenging activities indicated as percentage inhibition (Table 6). These data imply that methanolic extract of *Datura stramonium* is a potent scavenger of hydroxyl radical and may therefore prevent the various hydroxyl radical-induced pathological conditions. These powerful hydroxyl radical scavenging activities exhibited by *Datura stramonium* extract may be due to the presence of phytochemicals such as flavonoids and phenol as revealed by other data from the present study.

There is an established direct correlation between phenolic compounds and antioxidant activity. Phenolic compounds are generally

regarded as the most important antioxidative compound in the plant. As a result of this, the total antioxidant ability of a plant largely depends on its phenolic contents (Moskovitz, 2002). Data obtained in this study that *Datura stramonium* oil extract is rich in phenolic compounds. The concentration of these phenolic compounds increases as the concentration of the oil increases from 25 to 100%. The result is further confirmed by the data obtained from both GC-MS and HPLC analysis of the *Datura stramonium* oil extract. The high antioxidant activity of *Datura stramonium* oil and its rich phenol contents further confirm the strong correlation between antioxidant activities and the concentration of phenolic compounds.

Data obtained from GC-MS analysis of the oil extract showed that *Datura stramonium* oil extract contained phenolic compounds, fatty acids, aliphatic, amide, and alkene compounds. The presence of phenolic and other compounds in the oil extract further lend credence to the excellent antioxidant capacity of the oil extract and showed in the present study.

Furthermore, HPLC analysis confirmed the presence of catechin, rutin, and kaempferol in the *Datura stramonium* oil extract. Catechins are flavonols that belong to polyphenolic compounds. The complex of Catechin, epicatechin, and epigallocatechingallate, are the major flavonoids found in green tea. Data from many previous studies revealed that catechin plays a vital role in protection against degenerative diseases owing to its strong antioxidant power (Li et al. 2018). Many other researchers have also established an inverse relationship between catechin intake and the risk of cardiovascular diseases (Ikeda et al. 2018). Similarly, Cosarco et al. (2019) observed that catechin and its derivatives represent a class of phenolic compounds with a very wide therapeutic potential which includes; antiallergic properties, immunostimulatory, antidiabetic, antibacterial, antiviral, antioxidant, antitumor, lipid-lowering, and cardiovascular effects. Meanwhile, Rutin also referred to as rutoside, quercetin-3-rutinoside, and sophorin is a flavonol and is abundant in many plants, such as apples and tea. It is a very important nutritional component of food substances. Chemically rutin is a glycoside that is made up of

flavonolicaglyconequercetin and disaccharide rutinose. Previous studies have implicated rutin in several pharmacological activities such as antioxidant, cytoprotective, vasoprotective, anticarcinogenic, neuroprotective, and cardioprotective activities (Sharifi et al. 2018). Similarly, Kaempferol is a yellow compound and one of the most encountered aglycone flavonoids in the form of a glycoside. It is a tetrahydroxyflavone in which the four hydroxy groups are positioned on carbon 3, 5, 7, and 40. It is a natural flavonol-type flavonoid that can be found in various parts (fruits, leaves, fruits, and flowers) of plants such as tea and many common vegetables. Kaempferol and its glycosylated derivatives have been identified in different medicinal plants and have been shown to exhibit cardioprotective, neuroprotective, anti-inflammatory, antidiabetic, antioxidant, antimicrobial, antitumor, and anticancer properties (Sharifi et al. 2018).

CONCLUSION

In conclusion, our results indicated that *Datura stramonium* fruit oil possesses potent antioxidant potentials and this might be as a result of the presence of potent secondary metabolites as revealed by GC-MS and HPLC analysis of the fruit oil.

RECOMMENDATION

The authors recommend further studies, such as the toxicological potential of the plant to ascertain its suitability for human consumption or otherwise.

SOURCE OF FUNDING

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Adetola O, Alabi O, Abdulrauf I. (2016). Investigating storage duration and packaging materials on quality of fresh palm oil. *FJRS*. 12(2): 252-9.
- Ajayi GO, Olagunju JA, Ademuyiwa O, Martins OC. (2011). Gas chromatography-mass spectrometry

- analysis and phytochemical screening of ethanolic root extract of *Plumba gozeylanica*, Linn. *J. Med. Plant Res.* 5: 1756-61.
- Akinhanmi TF, Atasié VN. (2008). Chemical Composition and Physicochemical Properties of Cashew nut (*Anacardium occidentale*) Oil and Cashew nut Shell Liquid. *J. Agric. Food Environ. Sci.* 2: 1-10.
- AlKhattaf NF, Alraddadi AM, Aljarbou MA, Arnauti MA, Alfaleh AM, Hammouda SA. (2020). Determining the correlation between olive oil consumption, BMI, and waist circumference in the adult Saudi population. *J. Taibah Univ. Medical Sci.* 15(5): 380e386
- AOCS. (1990) Determination of saponification value. AOCS Publishing, Champaign, IL. USA.
- AOCS. (2004). Official methods and recommended practices of the American Oil Chemists' Society. AOCS Publishing, Champaign, IL. USA.
- AOCS. (2005). Official Methods of Analysis Cd 8b-90. AOCS Publishing, Champaign, IL. USA.
- Asari MA, Zulkaflee MH, Sirajudeen KNS, MohdYusof NA, MohdSairazi NS. (2019). Tualang honey and DHA-rich fish oil reduce the production of pro-inflammatory cytokines in the rat brain following exposure to chronic stress. *J. Taibah Univ. Medical Sci.* 14(4): 317e323
- Badmus JA, Adedosu OT, Fatoki JO, Adegbite VA, Adaramoye AO, Odunola OA. (2011). Lipid Peroxidation Inhibition and Antiradical Activities of Some Leaf Fractions of *Mangifera indica*, *Acta Pol. Pharm.* 68(1): 23-9.
- Bouziri A, Hamdi A, Borgi A, Hadj SB, Fitouri Z, Menif K, Jaballah NB. (2011). *Datura stramonium* L. poisoning in a geophagous child: A case report, *Int. J. Emerg. Med* 4:31.
- CODEX Alimentarius Commission. Codex standard for named vegetable oils-"Codex stan 210-1999". Food and Agricultural Organisation of the United Nations and the World Health Organisation. 2005.
- Cosarco S, Tunase C, Muntean DL. (2019). Therapeutic aspects of catechin and its derivatives – an update. *Acta Bio. Marisiensis* 2(1): 21-29.
- Dijkstra AJ. (2015). Vegetable Oils: Composition and Analysis: In Caballero B, Finglas P, Toldra F. (Eds): *Encyclopedia of Food and Health*. Academic Press Vol. 5 pp 4006.
- Elgohary HM, KH S, Jaouni A, Selim SA. (2018). Effect of ultrasound-enhanced *Nigella sativa* seeds oil on wound healing: An animal mode. *J. Taibah Univ. Medical Sci* 13(5): 438e443.
- Ganu GP, Jadhav SS, Deshpande AD. (2010). Antioxidant and antihyperglycemic potential of methanolic extract of bark of *mimusopselengi* l. In mice. *Int. J. Phytomedice* 2: 116-123.
- Halliwell B, Gutteridge J, Aruoma O. (1987). The deoxyribose method: A simple "test-tube" assay for determination of rate constants for reactions of hydroxyl radicals *Anal. Biochem* 165: 215.
- Halliwell B, Gutteridge JMC. (1999). *Free radicals in biology and medicine*, 3rd ed. Oxford: University Press, United Kingdom.
- Ikeda A, Iso H, Yamagishi K, Iwasaki, M Yamaji T, Miura T, Sawada N, Inoue M. Tsugane S, JPH Study Group. (2018). Plasma tea catechins and risk of cardiovascular disease in middle-aged Japanese subjects: The JPHC study. *Atherosclerosis*. 277: 90-97.
- Kumar AK, Ramachandra SS, Narsu L. (2010) Pharmacognostic and phytochemical investigations of roots of *Hibiscus micranthus* Linn. *Research Journal of pharmaceutical, Biological and Chemical Sciences*. 1(4): 324–337.
- Li F, Jin H, Xiao J, Yin X, Liu X, Li D, Huang Q. (2018). The simultaneous loading of catechin and quercetin on chitosan-based nanoparticles as effective antioxidant and antibacterial agent. *Food Res Int* 111: 351-60.
- McDonald S, Prenzler PD, Autolovich M, Robards K. (2001). Phenolic content and antioxidant activity of olive extracts. *Food Chem* 73: 73-84.
- Medda S, Fadda A, Dessena L Mulas M. (2021). Quantification of Total Phenols, Tannins, Anthocyanins Content in *Myrtus communis* L. and Antioxidant Activity Evaluation in Function of Plant Development Stages and Altitude of Origin Site. *Agronomy* 11(6): 1059.
- Mensor LL, Menezes FS, Leitao GG, Reis AS, dos Santos TC, Coube CS, Leitao SG. (2001). Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. *Phytother. Res* 15: 127.
- Moskovitz J, Yim MB, Chock PB. (2002). Free radicals and disease. *Arch BiochemBiophys* 397(2): 354-359.
- Mozaffarian D, Clarke R. (2009). Quantitative effects on cardiovascular risk factors and coronary heart disease risk of replacing partially hydrogenated vegetable oils with other fats and oils. *Eur J Clin Nutr* 63(S2): S22.

- Muhammad NO, Bamishaiye E, Bamishaiye OM, Usman L, Salawu M, Mikhail N, Oloyede H. (2011). Physicochemical Properties and Fatty Acid Composition of *Cyperus esculentus* (Tiger Nut) Tuber Oil. *Biores. Bull* 5: 51-4.
- Norton S. (2010). Toxic effects of plants; In Klaassen CD, Caserett D. (Eds): *Toxicology, the basic science of poisons*. McGraw Hill. 1110, New York Vol 7 pp 449-450.
- Oyaizu M. (1986). Studies on Products of Browning Reactions: Antioxidative Activities of Product of Browning Reaction Prepared from Glucosamine. *J-STAGE* 44: 307-315
- Parashuram M. (2011). Isolation of 11,12,13,17-Tetrahydroxy-(Hydroxymethyl)-10-Nitrodotriacontahydrospiro [Indeno[5,6-A]Hexacene-2,2 -Pyran]=3,6(1H,18bh) Dione and its spectroscopic characterization and biological activities of bimetals from seeds of *Datura stramonium*. *Asian J Bioch Pharm Res* 3(1): 501-506
- Pearson D. (1970). *The Chemical Analysis of Food*. 6th ed. J.A. Churchill, London.
- Ramadan MF, Sharanabasappa G, Seetharam YN, Seshagiri M, Moersel JT. (2006) Characterisation of fatty acids and bioactive compounds of kachnar (*Bauhinia purpurea* L.) seed oil. *Food Chem* 98(2): 359-65.
- Sanchez-Moreno C. (2002). Review: Methods Used to Evaluate the Free Radical Scavenging Activity in Foods and Biological Systems. *Food Sci. Technol. Int.* 8: 121.
- Sharifi-Rad M, Fokou PVT, Sharopov F, Martorell M, Ademiluyi OA, Rajkovic J, Salahi B, Martins N, Iriti M, Sharifi-Rad J. (2018). Antiulcer agents: From plant extracts to phytochemicals in healing promotion. *Molecules* 2018; 23(7): 1751.
- Szollosi R, Varga IS. (2002). Total antioxidant power in some species of *Labiatae* (Adaptation of FRAP method). *Acta Biol. Szeged.* 46(3-4): 125-7.
- Tsai JC, Huang GJ, Chiu TH, Huang SS, Huang SC, Huang TH, Lai SC, Lee VY. (2011). Antioxidant activities of phenolic components from various plants of *Desmodium species*. *Afr. J. Pharm. Pharmacol* 5: 468-476.