

Effects of Aqueous Leaf Extract of *Adansonia Digitata* (baobab) on Alloxan-induced Diabetes Mellitus on Endocrine Pancreas of Adult Male Wistar Rats

Journal of Experimental Research

September 2020, Vol 8 No 3

Email: editorinchief.erjournal@gmail.com
editorialsecretary.erjournal@gmail.com

Finbarrs-Bello E*¹, Neboh EE², Nnaji KS¹, Nwife GO³

¹Department of Anatomy, Faculty of Basic Medical Sciences. Enugu State University of Science and Technology, Parklane. Enugu State. Nigeria.

²Department of Medical Biochemistry, Faculty of Basic Medical Sciences. Enugu State University of Science and Technology, Parklane. Enugu State. Nigeria.

³Department of Anatomy, Faculty of Basic Medical Sciences. Alex Ekwueme University Udufu- Alike Ikwo Ebonyi State. Ebonyi State. Nigeria.

*Author for Correspondence: elizabeth.finbarrs-bello@esut.edu.ng

ABSTRACT

Adansonia digitata (Ad) also known as the baobab tree is characteristic of the Sahelian region and belongs to the Malvaceae family. Our study evaluated the possible therapeutic potential of *Adansonia digitata* leaf extract in the pancreas of alloxan-induced diabetic Wistar rats. Thirty-six (36) Wistar rats were randomly divided into 9 groups of four rats per group at the end of two weeks acclimatization. Group(g) 1(control group) received 0.1ml/kg/bw normal saline/14days, g2 (Diabetes negative control) received alloxan (150mg/kg)/2days, g3 (Low dose) received 150mg/kg/bw Alloxan + 200mg/kg/bw Ad extract g4 (Medium dose) received 150mg/kg/bw Alloxan +400mg/kg/bw Ad extract, g5 (High dose) received 150mg/kg/bw Alloxan +600mg/kg/bw Ad extract, g6 (Standard) received 150mg/kg/bw Alloxan + 150mg/kg/bw Metformin, g7 (Low dose) received 200mg/kg/bw Ad extract only, g8 (Medium dose) received 400mg/kg/bw Ad extract only and g9 (High dose) received 600mg/kg/bw Ad extract only. The animals were humanely sacrificed after 2 weeks of experiment under Ketamine anesthesia. Blood was collected via cardiac puncture for hormonal assay, after which a midline laparotomy was performed and the pancreas was harvested and fixed for histological studies. The study indicates that administration of Ad extracts reduced the weight of the animals in a dose-dependent manner, temporarily reduced blood glucose, increased glucagon and insulin concentrations, but was unable to restore the diabetic pancreas to its normal state. This may be attributed to the high carbohydrate content of the extract and absence of phytonutrients like tannin. The findings show that the leaf extract of Ad could be a potential and effective therapeutic target for short-term blood sugar regulation.

Keywords: Insulin, Glucagon, Diabetes Mellitus, Pancreas, Phytonutrients

INTRODUCTION

Medicinal plants have for thousands of years, been used as traditional treatment for numerous diseases and are generally considered to play a beneficial role in healthcare (Mohamed et al. 2015). *Adansonia digitata* (L.) called the baobab tree in both English and French is a characteristic of the Sahelian region (Savannah dry lands of sub-Saharan Africa) and belongs to the Malvaceae family (De Caluwe et al. 2010). With vast medicinal and non-medicinal benefits, local communities mainly utilize the leaves, pulp and seeds as a source of food and for income generation. It contains nutritionally significant levels of essential nutrients including fiber, proteins and minerals (Muthai et al. 2017). The fruit pulp is conventionally used against small pox, measles, diarrhea, scurvy, cough and

dysentery while the bark is used for fever treatment (Brendler et al. 2013).

The extract of the seed and fruit pulp of *Adansonia digitata* have been demonstrated to have anti-tumor action, as well as restore and modulate tumor markers levels (Elsaid, 2013). The fruit pulp extract also has a hepatoprotective influence on the liver (Mohammed et al, 2015). With its various parts being of utmost importance, Singh et al in his 2013 review stated that it possesses anti-oxidant, antipyretic, analgesic, antimicrobial, antiviral; as well as antidiabetic properties.

Diabetes mellitus on the other hand is a metabolic disorder that results from a reduction in insulin available for normal function of many cells in the body (Mohammed et al., 2015). This results from progressive destruction of Beta cells of the islets of Langerhans (Haligur, 2012). It is considered one of the main threats to

human health in the 21st century, and is associated with long-term complications that affect almost every part of the body; often leading to kidney failure, blindness, heart and blood vessel disease, stroke, amputations and nerve damage (Sournya and Srilatha, 2011). Although a chronic and progressive disease, initial lesions can be seen in very early stages, as also corroborated by previous studies where initial lesions were seen to occur after three days of diabetes mellitus induction in rats (Haligur, 2012). Clinically, the diagnosis of diabetes is based on a combination of hyperglycemia, variable loss of C-peptide secretion and dependence on exogenous insulin for survival, as well as destruction of pancreatic beta cells in autoimmune type 1 diabetes (Chitale, 2014). The pancreatic islets secrete hormones of vital importance, especially in the regulation of glycaemia, and blood glucose concentration. According to Saladin (2018), insulin insufficiency or inaction is well known as the cause of diabetes mellitus.

Alloxan is a toxic glucose analogue that preferentially accumulates in pancreatic beta cells, generating harmful hydroxyl radicals, as well as inhibiting beta cell glucose sensor or glucokinase (Lenzen, 2007). It is known to cause damage to the β -cells (general necrosis), leaving islets composed almost entirely of α -cells at approximately ninety-six hours after alloxan treatment. Hence, alloxan-induced diabetes is one of the most widely used and validated experimental model of chemical and drug-induced diabetic model for preclinical studies.

The present study investigated the possible therapeutic potential of the leaf extract of *Adansonia digitata* (baobab) on alloxan-induced diabetes mellitus in adult male Wistar rats.

MATERIALS AND METHODS

Collection, authentication and preparation of plant materials

Fresh leaves of *A. digitata* were procured from a local dealer in Kaduna state. It was identified by a botanist in the Department of Plant Sciences and Biotechnology, University of Nigeria, Nsukka. The extraction method described by Bello and Nwoso (2015)

was used. Five hundred (500g) of the powdered plant material was soaked in 1500ml of distilled water for 72 hours. Separation of the plant extract was carried out using Whatman No. 4 filter paper. The resulting extract was then concentrated using rotatory evaporator, and stored at 4°C before use.

Experimental animals and ethical approval

A total of 36 adult male wistar rats were purchased from the animal house of the University of Nigeria, Enugu Campus. The animals were bred, housed in netted iron cages and provided easy access to food (Grower's mash, from New market Enugu) and water ad libitum; maintained under standard laboratory conditions (temperature 20°C to 24°C, with relative humidity of 60-70% under 12 hours light and day cycles) and allowed to acclimatize for two weeks prior to the experiment in the Animal facility of Enugu State University of Science and Technology, (ESUT) College of Medicine, Enugu. At the end of acclimatization, the animals were randomly divided into nine (9) groups of four (4) animals each. The body weights of the animals were obtained before and after acclimatization, and also at weekly intervals during the experiment. The protocol for conducting the in vivo study in Wistar rats was approved by the Institutional Animal Ethical Committee (IAEC), Enugu State University of Science and Technology.

Induction of Diabetes using Alloxan

Alloxan Stock solution was prepared by dissolving alloxan monohydrate (Sigma-Aldrich, USA) (0.9g) in distilled water (6cm³), and diabetes was induced by single intraperitoneal injection of alloxan monohydrate (150mg/kg). The volume of the solution containing 150mg/kg given to each rat was determined by its weight. After a period of two days, the rats with blood glucose level greater than 200mg/dl was considered diabetic and used for the research work. The method of Mohammad and Hauwa'u (2013) was adopted in the study with slight modification.

Group 1(control group)- 0.1ml/kg/bw normal

saline/14days

Group 2 (Diabetes negative control) - Alloxan (150mg/kg)/2days

Group 3 (Low dose)-150mg/kg/bw Alloxan + 200mg/kg/bw *Adansonia digitata* extract

Group 4 (Medium dose)-150mg/kg/bw Alloxan +400mg/kg/bw *Adansonia digitata* extract

Group 5 (High dose)-150mg/kg/bw Alloxan +600mg/kg/bw *Adansonia digitata* extract

Group 6 (Standard)-150mg/kg/bw Alloxan + 150mg/kg/bw Metformin

Group 7 (Low dose)-200mg/kg/bw *Adansonia digitata* extract only

Group 8 (Medium dose)-400mg/kg/bw *Adansonia digitata* extract only

Group 9 (High dose)-600mg/kg/bw *Adansonia digitata* extract only

Determination of Blood Glucose Level

Blood samples of rats were collected by cutting the tail tip of the rats for glucose determination before administration of extract, after administration of the extract for 7 days and after administration of the extract for 14days. Determination of the blood glucose level was carried out using a glucometer (BG Check) and results were reported in mg/dl.

Hormonal assay

At the end of the experiment, blood samples were collected via cardiac puncture from all rats. Serum was then separated and stored at -20°C until the hormonal assay. Serum levels of total insulin and glucagon were determined by **ELISA (Enzyme-linked Immune-sorbent Assay) by Micallef et al.(1995)**. The desired number of coated wells was properly secured in the holder and 100µl of standards, specimens, and controls was dispensed into appropriate wells. It was thoroughly mix for 30seconds and incubated at room temperature (18-25°C) for 60 minutes. The incubated mix was removed by flicking plate contents into a waste container. The microtiter well was rinse and flick 5 times with distilled or deionized water. The wells was stroked sharply onto absorbent paper or towels to remove all residual water droplets, then 100µl of TMB reagent was dispense into each well and mixed gently for 10 seconds. The for

reaction was finally stopped by the addition of 100µl of stop solution to each well and mixed gently for another 30seconds. The absorbance was then read at 450nm with a microtiter well reader within 15 minutes

Histopathology studies

At the end of the study, the animals were humanely sacrificed under anesthesia (ketamine hydrochloride), twenty-four hours after the last treatment. Sections of the pancreas were collected for histopathological examination. The samples were fixed in 10% phosphate buffered formalin for a minimum of 48 hours. The tissues were subsequently trimmed, dehydrated in 4 grades of alcohol (70%, 80%, 90% and absolute alcohol), cleared in 3 grades of xylene and embedded in molten wax. On solidifying, the blocks were sectioned, 5µm thick with a rotary microtome, floated in water bath and incubated at 60°C for 30 minutes. The 5µm thick sectioned tissues were subsequently cleared in 3 grades of xylene and rehydrated in 3 grades of alcohol (90%, 80% and 70%).The sections were then stained with Hematoxylin for 15 minutes. Blueing was done with ammonium chloride. Differentiation was done with 1% acid alcohol before counter staining with Eosin. Permanent mounts were made on degreased glass slides using a mountant; DPX.

The prepared slides were examined with a Motic™ compound light microscope (Motic BA410E Elite Research Compound Microscope; Motic Asia, Hong Kong) using x 4, x 10 and x 40 objective lenses. The photomicrographs were taken using a Motic™ 5.0 megapixels microscope camera at x 160 magnification

Statistical analysis

Data obtained was expressed as the mean ± standard deviation. They were analyzed using Statistical Package for Social Sciences (SPSS version 21; IBM SPSS, Chicago, Illinois, USA) software package. Comparisons within and between groups were done using one way ANOVA with Tukey post hoc test. The level of significance was set to p<0.05.

RESULTS

Phytochemical analysis

The phytochemical analysis result shows that the aqueous leaf extract of *Adansonia digitata* used in the experiment contains alkaloids, flavonoids, glycosides, saponin and terpenoids. The carbohydrate content of the extract was very high, followed by protein, lipids and crude fibre respectively (table 1).

Table 1: Proximate Analysis of *Adansonia digitata* leaves extract

Parameters	Values in %
Moisture	0.81
Protein	14.35
Crude fibre	7.22
Ash	1.43
Fats and Oil	10.00
Carbohydrate	66.19

Table 1: The phytochemical analysis result shows that the aqueous leaf extract of *Adansonia digitata* Moisture and ash content are negligible.

Body weight

From table 2A below our results (values without superscripts) showed that there was no significant difference in weight among and between groups from day one to day 15 (after acclimatization). As revealed in table 2B, there was an increase (19.2%) in body weight of group 1 at the end of the experiment. In contrast, group 2 animals that served as the positive control (untreated diabetic group) showed a 10.7% decrease in body weight at the end of the experiment. Among the diabetes-treated groups (3, 4, 5 and 6) which were given low, medium, high doses of the aqueous leaf extract of *Adansonia digitata* and Metformin respectively after alloxan induction, only group 3 showed an increase in body weight (27.5%) while groups 4, 5 and 6 showed a decrease (8.24%, 10.62% and 3.97% respectively) in their body weight at the end of the experiment. Group 7 and 8 showed a 4.98% and 5.48% decrease in their body weight respectively while group 9 also had a 5.26% decrease in body weight at the end of the experiment.

Table 2A: Results of Mean± SD of the body weights in gram of all the experimental animals

Groups	Day1	Day 8	Day 15
1	120.0±9.13	127.5±5.00	143.1±13.07
2	141.7±2.87	139.2±13.21	126.5±10.62*
3	152.5±6.46	174.7±16.07	194.5±15.98*
4	178.3±14.43*	139.8±9.20	163.6±4.01 ^β
5	171.3±27.80*	178.2±73.57	153.1±10.57 ^{βc}
6	173.8±15.48*	154.9±29.48	166.9±29.30 ^β
7	168.6±17.05*	161.2±14.61	160.2±15.83 ^β
8	153.2±10.19	144.9±11.30	144.8±11.95 ^{βc}
9	140.7±16.0	143.5±12.69	148.1±13.71 ^{βc}

* = p<0.05 compared with the control group 1; ^β =p<0.05 compared with the control group 2; ^{CDEFGHI} =p<0.05 compared with the control group 3,4,5,6,7,8,9.

Table 2B: Weekly and percentage changes in body weights (g)

Groups	Week 1 Changes (d8 - d1)	Week 2 Changes (d15 - d1)	% changes (d1 - d8)	% changes (d8-d15)	% total changes (d15 - d1)
1	7.5±9.13	23.1±3.94	6.5	12.24	19.2
2	2.5±10.34	15.2±7.75	1.7	1.26	10.7
3	22.2±9.61	42.0±9.52	14.56	11.33	27.5
4	38.5±5.23	14.7±10.42	21.59	17.02	8.24
5	7.2±5.23	18.2±17.23	4.2	14.09	10.62
6	18.9±14.0	6.9±13.82	10.87	7.75	3.97
7	7.4±2.44	8.4±1.22	4.39	0.65	4.98
8	8.3±1.11	8.4±1.1	5.42	0.06	5.48
9	2.8±3.31	7.4±3.31	6.5	12.24	5.26

Data analyzed at Mean± SD and significance at p<0.05

Fasting blood glucose

This table shows the effect of the extract on fasting blood glucose levels at weekly intervals. On Day 1, there is a statistically significant (p<0.05) increase in group 2 glucose level (422.7 ± 135.9 mg/dl) and a significant (p<0.05) decrease in group 7 when compared to the normal control (group 1), as well as statistically significant decreases in groups 3, 5, 7, 8 and 9, compared to the diabetic control group (group 2). At the end of the first week of extract administration (Day 8), there shows a decrease in fasting blood glucose levels among the diabetes induced groups (groups 3, 4 and 6) compared to day 1. Also, there are increases in glucose level among groups 1, 2, 5, 7, 8 and 9 with the diabetic group (group 2) showing the highest level of glucose concentration (439.2 ± 13.21 mg/dl) compared to day 1. However, none of these are statistically significant difference (p>0.05)

except groups 6 and 7 which shows a significant decrease (p<0.05) when compared to group 2. By the second week of the experiment, there is an increase in fasting blood glucose levels among all the groups when compared to the previous week (Day 8), but a decrease in the group 6 compared to Day 1. Group 2 shows the highest concentration (466.5 ± 110.62 mg/dl) followed by group 6 (309.8 ± 111.0 mg/dl) which was statistically significant (p<0.05) when compared to group 1. There were also statistically significant differences (p<0.05) among groups 3-6 showing decrease in glucose concentration compared to group 2, and also among groups 7-9 when compared to group 6. The change in glucose concentration appears prominently in group 5 (69.9 %) and least in group 2 (10.4 %) at the end of the experimental period of two weeks.

Table 3: Results of the effects of aqueous extract of *Adansonia digitata* leaf on fasting blood glucose level (mg/dl)

Group	Day 1	Day 8	Day15	Changes (D15-D1)	% Changes
1	85.5±3.7	95.2±6.8	110.0±4.32	24.5±0.62	28.7
2	422.7±135.9*	439.2±13.21	466.5±110.62*	43.8±25.3	10.4
3	181.8±71.3 ^β	153.8±92.1	223.5±68.4 ^β	41.8±2.9	22.9
4	191.3±134.1	120.33±69.2	217.3±142.6 ^β	26.0±8.0	13.6
5	149.8±64.3 ^β	199.0±200.95	254.5±117.89 ^β	104.8±53.6	69.9
6	389.5±185.2	198.8±77.5 ^β	309.8±111.0* ^β	63.8±74.2	17.1
7	76.3±4.3* ^β	84.3±5.50 ^β	108.3±47.5 ^f	37.5±43.2	49.4
8	80.5±5.5 ^β	92.5±13.8	113.8±78.9 ^f	33.3±73.5	41.4
9	76.5±1.3 ^β	82.0±2.5	120.0±30.0 ^f	43.5±28.7	63.4

*= p<0.05 compared with the control group 1; ^β= p<0.05 compared with the control group 2; ^{CDEFGHI}= p<0.05 compared with the control group 3,4,5,6,7,8,9.

Hormonal analysis

From the table, there were significant increases in all the groups compared to the negative control (group 1), with the exception of group 2 (untreated diabetic group) which has the least insulin concentration at the end of the experiment (9.86 ± 0.87 mIU/mL). The increase in insulin concentration was more pronounced in group 5 (22.12 ± 3.94 mIU/mL), followed by groups 6, 4 and 7 respectively. The differences in groups 3-9 were statistically significant (p<0.05) when compared to group 2. Likewise, there are statistically significant differences (p<0.05) between groups 2, 4, 5 and

6 when compared to the control group. The glucagon concentrations decreased significantly (p<0.05) at the end of the experiment among groups 2-9 when compared with the negative control group (group 1), while group 2 has the lowest glucagon concentration (25.44 ± 1.36 pg/ml). There are also significant increases in glucagon among the groups, with group 4 having the highest glucagon concentration (63.86 ± 18.41 pg/ml), followed by group 5, 8, 9 and 7 respectively. The differences in groups 4 and 5 were statistically significant (p<0.05) compared to group 2.

Table 4: Results of the effect of aqueous leaf extract of *Adansonia digitata* on hormonal (insulin & glucagon) level in the Wistar rats

Groups	Insulin concentration (mIU/mL)	Glucagon concentration (pg/ml)
1	15.86±2.19	107.08±6.53
2	9.86±0.87* ^{cdefgh1}	25.44±1.36*
3	20.66±1.13 ^β	48.65±27.99*
4	21.39±2.24* ^β	63.86±18.41* ^β
5	22.21±3.94* ^β	55.38±7.09* ^β
6	21.76±2.05* ^β	39.33±4.49*
7	20.84±1.35 ^β	50.41±6.11*
8	17.41±2.64 ^β	51.34±2.75*
9	20.19±1.01 ^β	50.49±2.06*

*=p<0.05 compared with the control group 1; ^β =p<0.05 compared with the control group 2; ^{CDEFGHI}=p<0.05 compared with the control group 3,4,5,6,7,8,9.

Histological findings

The endocrine pancreas showed normal histoarchitecture in group 1 (negative control) while group 2 (positive control) showed decreased number of islets, confirming diabetes. Groups 3, 4 and 5 (extract-treated diabetic groups) showed decreased islet numbers and sizes while group 6 (metformin-

treated diabetic group) showed regeneration by increased number of islets. Groups 7, 8 and 9 (extract-treated non-diabetic groups) showed normal histoarchitecture of the endocrine pancreas. These effects of *Adansonia digitata* aqueous leaf extract on the histology of the endocrine pancreas are presented in the photomicrographs below.




Plate 1: Sections of the pancreas collected from the animals in this group (group 1, normal saline) showed the normal histo-architecture of both the exocrine and endocrine pancreas. The pancreatic lobules contained normal sized pancreatic islets (arrow) which are composed of a cluster of pale eosinophilic cells with eccentrically located nuclei and abundant cytoplasm. Exocrine pancreas (EP).H&E. x160.




Plate 2: Sections of the pancreas collected from the animals in this group (group 2, alloxan monohydrate 150mg/kg) showed normal histo-architecture of the exocrine pancreas. However, the endocrine pancreas showed a marked decrease in the number and sizes of the pancreatic islets. The few observed pancreatic islets (arrow) were scanty, composed of tightly packed islet cells and are relatively inconspicuous. Exocrine pancreas (EP). H&E.x160




Plate 3: sections of the pancreas collected from the animals in this group (group 3, alloxan monohydrate 150mg/kg + *Adansonia digitata* leaf extract 200mg/kg)

showed normal histo-architecture of the exocrine pancreas. the endocrine pancreas showed a marked decrease in the number and sizes of the pancreatic islets (arrow). the few observed pancreatic islets (arrow) were relatively inconspicuous. exocrine pancreas (ep); blood vessel (bv); pancreatic duct (pd). H&E. x160.




Plate 4: Sections of the pancreas collected from the animals in this group (group 4, alloxan monohydrate 150mg/kg + *Adansonia digitata* leaf extract 400mg/kg) showed normal histo-architecture of the exocrine pancreas. The endocrine pancreas showed a marked decrease in the number and sizes of the pancreatic islets (arrow). The few observed pancreatic islets (arrow) were relatively inconspicuous. Exocrine pancreas (EP); Pancreatic duct (PD); Blood vessel (BV). H&E.x160.




Plate 5: Sections of the pancreas collected from the animals in this group (group 5, alloxan monohydrate 150mg/kg + *Adansonia digitata* leaf extract 600mg/kg) showed normal histo-architecture of the exocrine pancreas. The endocrine pancreas showed a marked decrease in the number and sizes of the pancreatic islets (arrow). The few observed pancreatic islets (arrow) were relatively inconspicuous. H&E. x160.




Plate 6: Sections of the pancreas presented in this group (group 6, alloxan monohydrate 150mg/kg + metformin 150mg/kg) showed a marked decrease in the number of the pancreatic islets. A few of the pancreatic islets (arrow) showed marked increase in the number and size of the pancreatic islet cell. This is highly suggestive of pancreatic islet regeneration. Exocrine pancreas (EP).H&E.x160.




Plate 7: Sections of the pancreas collected from the animals in this group (group 7, *Adansonia digitata* leaf extract 200mg/kg) showed the normal histo-architecture of both the exocrine and endocrine pancreas. The pancreatic lobules contained normal sized pancreatic islets (arrow) which are composed of a cluster of pale eosinophilic cells with eccentrically located nuclei and abundant cytoplasm. Exocrine pancreas (EP).H&E. x160.




Plate 8: Sections of the pancreas collected from the animals in this group (group 8, *Adansonia digitata* leaf extract 400mg/kg) showed the normal histo-architecture of both the exocrine and endocrine pancreas. The

pancreatic lobules contained normal sized pancreatic islets (arrow) which are composed of a cluster of pale eosinophilic cells with eccentrically located nuclei and abundant cytoplasm. Exocrine pancreas (EP).H&E.x160.




Plate 9: Sections of the pancreas collected from the animals in this group (group 9, *Adansonia digitata* leaf extract 600mg/kg) showed the normal histo-architecture of both the exocrine and endocrine pancreas. The pancreatic lobules contained normal sized pancreatic islets (arrow) which are composed of a cluster of pale eosinophilic cells with eccentrically located nuclei and abundant cytoplasm. Exocrine pancreas (EP).H&E.x160.

DISCUSSION

In this study, the use of the leaf extract of *Adansonia digitata* on alloxan-induced diabetic rats was to investigate the relationship between *A. digitata* leaves extract and some of the diabetic symptoms with an aim to establishing its usefulness in the treatment of diabetes. As stated by Melmed et al. (2010), blood plasma glucose concentration is one of the criteria for assessing for diabetes. Diabetes mellitus is marked by increased blood glucose sequel to decreased insulin secretion by β -cells of the pancreas (Rang et al. 2012), as observed in the untreated diabetic control (group 2). Hyperglycemia therefore, is a primary factor in the development of diabetes complications, and decreases in average blood glucose have a profound effect to prevent complications in diabetes mellitus (Diabetes Control and Complications Trial Research Group, 1993). The fasting blood glucose profile shows that *A. digitata* leaves extract was capable of lowering

plasma glucose levels, as seen in the diabetic groups treated with low and medium doses of the extract (200mg/kg and 400mg/kg respectively) after the first week of the experiment. This supports previous studies on other parts of the *Adansonia digitata* plant which were shown to possess antidiabetic potential (Mohammad and Hauwa'u, 2013). The blood glucose levels of the diabetic group treated with a high dose of the extract (600mg/kg), as well as the normal groups treated with different doses of the extract (groups 7-9), on the other hand, were increased after administration. However, at the end of the second week of the experiment, the blood glucose levels increased in all the diabetic and non-diabetic groups treated with the extract. This increase in blood glucose levels possibly could be as a result of the absence of tannins in the leaf extract, which was otherwise detected in the fruit pulp (Mohammad and Hauwa'u, 2013) and stem bark extracts of *A. digitata*. Tannins are however known for inducing regeneration of β -cells (Anderson and Polansky, 2002) and also as anti-hyperglycemic agents in diabetic rats (Pinent et al. 2004). More so, this increase in blood glucose might also have arisen owing to the high carbohydrate content of the extract, as Eizirik and Cnop (2010) reported that carbohydrate consumption increases demand on the β -cell for insulin secretion, which may lead to endoplasmic reticulum stress, as well as oxidative stress (Robertson et al. 2004) both of which can result in β -cell damage after a long period of time. It is also possible that a high-carbohydrate/lower-fat diet such as contained in the extract, on a prolonged period of time could increase insulin sensitivity and lower fasting glucose levels, as reported by Gower et al (2012). The increase in blood glucose levels after a decrease in the previous week could also have resulted from an increase in glucagon and insulin in the regulation of homeostasis, released in a pulsatile manner to regulate glucose production and metabolism (Farhy and McCall, 2009), in which glucagon promotes glycogenolysis and gluconeogenesis to increase endogenous blood glucose levels during prolonged fasting. It was also observed, however, that the blood glucose levels of the

diabetic group treated with the extract were significantly lower than that of the untreated diabetic group as well as the group treated with the standard drug (metformin). This suggests that the extract has an antagonistic effect between its high carbohydrate content and its hypoglycemic property via unclear mechanisms.

From the results, the insulin concentrations of all the groups were significantly elevated except in the diabetic control group which has the least concentration of insulin when compared with the negative control (group 1). This reduction in the insulin concentration is due to the selective destruction or necrosis of the insulin-producing β -cells of the pancreatic islets; thus causing diabetes. Insulin secretion is therefore inhibited due to the β -cell toxicity, As noted by Szkudelski (2001), the mechanism of cytotoxic action of alloxan on β -cells involve oxidation of essential sulphhydryl (-SH) group, inhibition of glucokinase, generation of toxic free radicals and disturbance in intracellular homeostasis with the resulting damage resulting in decreased insulin release and attendant hyperglycemia. Conversely, the increase in insulin was most pronounced in the diabetic rats treated with high dose of the extract (group 5), followed by the diabetic rats treated with metformin (group 6). This suggests that the insulin-increasing effect of the extract was dose-dependent, and more potent than that of metformin. The significant increase in insulin secretion can be attributed to a stimulation of the β -cells by flavonoids present in the extract (Benhabyles et al. 2015).

Insulin is known to regulate glucagon, as well as various β -cell signals that provide an inhibitory stimulus to the α -cells and suppress glucagon (Gromeda et al. 2007). Therefore, an increase in insulin secretion suppresses glucagon secretion and a decrease in insulin, in concert with a low glucose concentration, stimulates glucagon secretion (Cooperberg and Cryer, 2010). The findings of our study showed a significant decrease in the glucagon concentrations in all the groups compared to the negative control (group 1), as well as a significant increase in the diabetic rats treated with medium and high doses of the extract

(groups 4 and 5) compared to the positive control (group 2). This also confirms that the extract has a dose-dependent effect on pancreatic hormonal concentrations.

The histopathological observations in the present work on one hand showed that the β -cells in the untreated diabetic control group were significantly reduced in size and number, thus confirming diabetes. The decrease in islet size and β -cell number, as well as the histoarchitecture of the islets, is in agreement with Elkotby et al (2018). Therefore, the serum insulin level was decreased and the glucose concentrations were increased as shown in tables 3 and 4. The diabetic groups treated with varying doses of the extract (groups 3-5) all show marked decrease in size and number of β -cells, though this is more prominent in those treated with medium and high doses of the extract. These tally with the increased blood glucose levels in the groups. Thus, we can agree Karadimos et al. (2012) that the quantity of remaining β -cells was not sufficient to control blood glucose. On the other hand, the metformin treated diabetic group (group 6) shows the same symptoms as groups 3-5, but with a few of the islets showing an increase in the β -cells. This confirms metformin's ability to regenerate cells (El-Soud et al. 2016). The non-diabetic rats treated with different doses of the extract (groups 7-9) shows normal histoarchitecture of the pancreatic islets. This suggests that, while the high carbohydrate content may cause an increase in glucose concentrations, it is not enough to cause hyperglycemia ($>200\text{mg/dl}$) which is a characteristic of diabetes. Therefore, the extract has no adverse effects on normal rats, could lower blood glucose levels but on prolonged administration, might cause hyperglycemia in medium and high doses more than low doses, and subsequently lead to the onset of diabetes. In comparison, metformin increased the blood glucose levels even more than that of the extract, but also shows its curative potential in the regeneration of β -cells. The reasons for inducing hyperglycemia in diabetes treatment by the extract; after showing its hypoglycemic potential is currently unknown. This could be due to paucity of data as *Adansonia digitata* leaves have been poorly

investigated (Sokeng et al. 2019), in relation to diabetes.

The percentage decrease in body weight noticed in the diabetic group may be due to the adverse effects of diabetes to the body. This is in agreement with Eghoi and Paul (2016), who stated that diabetes is often associated with a characteristic loss of body weight, partially due to increased muscle wasting. Groups 7, 8 and 9 showed decreases in their body weights at the end of the experiment. This suggests that the extract does not cause weight gain irrespective of the dose of the extract given, thus it is not dose-dependent. However, comparing the percentage changes in body weight of animals in group 7, 8 and 9 to group 1, it can be suggested that the aqueous leaf extract of *Adansonia digitata* had no benefiting effect on the body weight of the animals. These observations agree with previous studies of Eghoi and Paul (2016). Among the diabetes-treated groups (3, 4, 5 and 6) which were given low, medium, high doses of the aqueous leaf extract of *Adansonia digitata* and metformin respectively after alloxan induction, only group 3 showed an increase in body weight while groups 4, 5 and 6 showed decreases in their body weight at the end of the experiment. This reduction in weight may be due to the presence of alkaloids in the extract (Burkill, 1994). Since the animals in this group (4, 5 and 6) also had increased blood glucose levels (13.6%, 69.9% and 17.1% respectively) as seen in table 3, it may also be suggested that they were also experiencing the weight-decreasing effect of diabetes as also noticed in the group 2. This also agrees to the idea that the aqueous leaf extract of *Adansonia digitata* was not able to maintain reduced blood glucose levels at the end of the experiment even when there was a decrease in the blood glucose levels in these groups during the first week (Table 4.5). Loss of body weight is a major consequence of diabetes in rats (Ramachandran et al. 2012). This could be due to dehydration and catabolism of fats and proteins (Rajkumar et al. 1991). The prevention of loss in body weight by the low extract dosage may be due to increasing glucose uptake in peripheral tissues or inhibition of catabolism by good glycemic control (Ambika et al. 2013).

To the best of our knowledge *Adansonia digitata* leaf extract therefore causes weight loss which may be due to the presence of alkaloids in the extract or simply as a result of the side effects of diabetes. There was a high increase in blood sugar levels in the diabetes-treated groups as well as a decrease in the pancreatic islets, which may be due to the high carbohydrate content of the extract. In contrast, the histoarchitecture of the non-diabetic groups treated with the extract remained normal while the blood sugar levels were minimally increased. This suggests that the extract has no adverse side effects on normal rats and so can possibly be effective as a temporal (not on prolonged basis) therapeutic measure against DM. The hormonal profiles which showed an increase in insulin concentration in all the groups may be due to the presence of flavonoids in the extract. Alternatively, the general decrease in glucagon concentrations may be as a result of the increase in insulin secretion.

CONCLUSION

In conclusion, the present study indicates that administration of *Adansonia digitata* extracts to rats with alloxan-induced diabetes, reduced the weight of the animals in a dose-dependent manner, temporarily reduced blood glucose, increased glucagon and insulin concentrations, but was unable to restore the diabetic pancreas to its normal state. The mechanism of this effect is yet unknown but is apparently related to the high carbohydrate content of the extract and absence of phytonutrients like tannin.

REFERENCES

- Ambika SR, Saravanan R, Thirumavalavan K. (2013). Antidiabetic and antihyperlipidemic effect of p-hydroxycinnamic acid on streptozotocin-induced diabetic Wistar rats. *Biomed. Aging Pathol.* 3: 253-257
- Anderson RA, Polansky MM. (2002). Tea enhances insulin activity. *J Agric. Food Chem.* 50: 7182-7186
- Bello OK, Nwoso SO. (2015). Radioactive potentials of methanolic/aqueous extract of *Adansonia digitata* and *Cochorouso litorious* leaf plant on gamma irradiated male wistar rats. *Journal of Natural Sciences Research.* 5(11): 81-93
- Benhabyles N, Arab K, Bouchenak O, Baz A. (2015). Phytochemical screening, hypoglycemic and antihyperglycemic effect of flavonoids from the leaves of Algerian *Olea europaea* L. in normal and alloxan-induced diabetic rats. *International Journal of Pharmacology.* 11;(5): 477-483
- Brendler T, Gruenwald J, Jaenicke C. (2013). Herbal remedies. Medpharm Scientific Publishers: Stuttgart, Germany
- Burkill HM. (1994). The useful plants of west tropical Africa. Royal Botanical Gardens *Kew, UK.*
- Buse JB, Rosenltock J, Sesti G. (2009). Liraglutide once a day versus Exenatide twice a day for type 2 diabetes: a 26-week randomized, parallel-group, multinational, open-label trial (LEAD-6). *The Lancet.* 374;(9683): 39-47
- Chitale AR. (2014). Morphology of Pancreatic Islets and Pathology of Pancreas in Diabetes Mellitus. In: RSSDI Textbook of Diabetes Mellitus [3rd ed.] by Humraj BC, Gumpeny RS, Ashok KD, Sri VM, Viswanathan M, Paturi VR [eds]. Jaypee Brothers Medical Publishers (P) Ltd: New Delhi, India; pp.45
- Cooperberg BA, Cryer PE. (2010). Insulin reciprocally regulates glucagon secretion in humans. *Diabetes.* 59;(11): 2936-2940.
- De Caluwe E, Halamora K, Van Damme P. (2010). *Adansonia digitata* L. – A review of traditional uses, phytochemistry and pharmacology. *Africa Focus.* 23: 11-51.
- Eghoi AM, Paul CW. (2016). The effect of methanolic leaf extract of *Adansonia digitata* on the histology of Wistar rats testes. *Advances in Biological Research.* 10;(6): 365-373.
- Eizirik DL, Cnop M. (2010). ER stress in pancreatic beta cells: The thin red line between adaptation and failure. *Sci. Signal.* 3(110): e7.
- Elkotby D, Hassan AK, Emad R, Bahgat I. (2018). Histological changes in islets of Langerhans of pancreas in alloxan-induced diabetic rats following Egyptian honey bee venom treatments. *Int. J Pure Appl. Zool.* 6(1): 1-6.
- Elsaid FG. (2013). The effects of seeds and fruit pulp of *Adansonia digitata* on Ehrlich Ascites carcinoma. *Food and Nutrition Sciences.* 4: 38-46.
- El-Soud NHA, El-Laithy NA, Mohamed NA, Youness ER, Wasseif ME, Yassen NN, Abdel-Latif AM. (2016). Honey versus metformin: Effects on pancreatic beta-cells in streptozotocin induced

- diabetic rats. *DerPharma. Chemica.* 8(16): 29-39
- Farhy LS, McCall AL. (2009). Pancreatic network control of glucagon secretion and counter-regulation. *Methods Enzymol.* 467: 547-581
- Gower BA, Goree LL, Chandler-Laney PC, Ellis AC, Casazza K, Granger WM. (2012). A higher-carbohydrate, lower-fat diet reduces fasting glucose concentration and improves β -cell function in individuals with impaired fasting glucose. *Metabolism.* 61(3): 358-365.
- Gromeda J, Franklin I, Wollheim CB. (2007). A-cells of the endocrine pancreas: 35 years of research but the enigma remains. *Endocrine Reviews.* 28(1): 84-116.
- Haligur M, Topsakal S, Ozmen O. (2012). Early degenerative effects of diabetes mellitus on pancreas, liver and kidney in rats: An immunohistochemical study. *Experimental Diabetes Research.* 120645
- Karadimos MJ, Kapoor A, Khattabo IEL, Sharma A. (2012). β -cell preservation and regeneration for diabetes treatment: where are we now? *Diabetes Manag. (Lond);* 2(3): 213-222.
- Lenzen S. (2007). The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia.* 51: 216-226.
- Melmed S, Polonsky KS, Larson PR, Kronenberg HM. (2010). *Williams Textbook of Endocrinology* (12th ed.). Elsevier Saunders: London; p. 1442-1447.
- Mohammad YG, Hauwa'u YB (2013). Hypoglycemic activity of methanolic fruit pulp extract of *Adansonia digitata* on blood glucose levels of alloxan induced diabetic rats. *Int. J Anim. Veter. Adv.* 5(3): 108-113
- Mohammed MA, Tajeldin IM, Mohammed AH, Hozeifa HM. (2015). Hepatoprotective effect of *Adansonia digitata* L. (Baobab) fruit pulp extract on CCL₄-induced hepatotoxicity in rats. *World Journal of Pharmaceutical Research.* 4(8): 368-377
- Muthai KU, Kaori MS, Muchugi A, Indieka AS, Demebele C, Mng'omba S. Jamnadass R. (2017). Nutritional variation in baobab (*Adansonia digitata* L.) fruit pulp and seeds based on geographical regions. *Food Sci. Nutr.* 5:1116-1129
- Pinent M, Blay M, Blade MC, Salvado MJ, Avola L, Ardevol A. (2004). Grape seed derived procyanidins have an anti-hyperglycemic effect in streptozotocin-induced diabetic rats and insulinomimetic activity in insulin-sensitive cell lines. *Endocrinology.* 145: 4985-4990
- Rajkumar L, Govindarajula P, Srinivasan N, Balasubramian K. (1991). Increased degradation of dermal collagen in diabetic rats. *Indian J Exp. Biol.* 29: 1081-1083
- Ramachandran S, Naveen KR, Rajinikanth B, Akbar M, Rajasekaran A. (2012). Antidiabetic, antihyperlipidemic and in vivo antioxidant potential of aqueous extract of *Anogiessus latifolia* bark in type 2 diabetic rats. *Asian Pac. J Trop. Dis.* 23: S596-S602.
- Rang HP, Dale MM, Ritter JM, Moore PK. (2012). *Pharmacology.* Churchill Livingstone: London. pp: 310.
- Robertson RP, Harmon J, Tran POT. (2004). Beta-cell glucose toxicity, lipotoxicity, and chronic oxidative stress in type 2 diabetes. *Diabetes.* 53;(suppl 1): 5119-5124
- Saladin KS. (2018). *Anatomy and Physiology: The Unity of Form and Function* [8th ed.]. McGraw-Hill Education: New York; 627-643
- Singh S, Varsha P, Shashi R. (2013). Medicinal uses of *Adansonia digitata* L: An endangered tree species. *JPSI.* 2;(3): 14-16
- Sokeng AJT, Sobolev AP, DiLorenzo A, Xiao J, Mannina L, Capitani D, Daglia M. (2019). Metabolite characterization of powdered fruits and leaves from *Adansonia digitata* L. (Baobab): A multi-methodological approach. *Food Chem.* 272: 93-108.
- Sournya D, Srilatha B. (2011). Late stage complications of diabetes and insulin resistance. *J Diabetes Metab.* 2;(9): 1-7.
- Szkudelski T. (2007). The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physiol. Res.* 50;(6): 536-46.