



ORGANOSOMATIC AND HAEMATOLOGICAL INDEX OF ALBINO WISTAR RATS (*Rattus norvegicus*) FED WITH CHICKEN FLAVOUR INDOMIE SEASONING

Opara, Kenneth¹, Oboho, Diligent^{*}, Nelson, Akwaowo²,
Udoigung, Nsima¹ Esigbone, Alero¹ and Edeke, Affiong³

¹Department of Animal and Environmental Biology, University of Uyo; Uyo
Nigeria

²Department of Animal Science, Federal Polytechnic Ugep Cross River State,
Nigeria

³Department of Chemical Sciences, Topfaith University Mkpatak, Akwa Ibom
State Nigeria

*Author for Correspondence: diligentoboho@uniuyo.edu.ng

ABSTRACT

The effects of the indomie seasoning (chicken flavour) on the organosomatic and haematological indices of Albino rats were investigated under laboratory conditions of sixteen (16) albino rats grouped into four (A-D) groups. Each group had 4 albino rats, treated groups were labelled A-C and the control was labelled D. The group means \pm SEM for each parameter was calculated SPSS (26.0). Means were separated using Turkey's post hoc test at 5% level of significance. The results showed that there was steady increase in the weight of the animals fed orally with 25mg/kg, 50mg/kg and 75mg/kg of the indomie seasoning a week(s) increases. It is also observed that there was a significant increase ($p > 0.05$) in the organ weight of the liver (hepatosomatic index) in all the groups compared to control. Also, the renosomatic index was significantly increased as the weeks increased. Results also showed that the indomie seasoning produces a significant change in the platelets counts when compared with the control. However, the result showed a general change in the haematopoietic counts of the albino rats. Since the effects of the seasoning on the general well-being of humans had been documented, it is therefore, pertinent for us to reduce the intake of the seasoning especially at low amount.

Keywords: Albino wistar rats, Haematological parameters, Indomie seasoning, Organosomatic index

INTRODUCTION

Depending on the brand, instant noodles are precooked, dry noodles made from wheat flour that are packaged with oil, seasoning, flavouring powder, and other components. (Amin and others, 2010). These noodles still exist today in many types of instant noodles, and some of its ingredients include refined flour made from weed, salt, and oil (Kim, 1996a). The great majority of supplements have been shown to work remarkably well with noodles. Most people call them junk food (Hope, 2009). According to Joseph et al. (2017), a specific meal consisting of these instant noodles is stated to be poor in vital nutrients including vitamins, protein, fiber, and minerals and high in added sugar, sodium, and other chemicals. These noodles are particularly popular because they are readily available, inexpensive, and simple to prepare—they only take two to three minutes. In recent times, there have been multiple brands of instant noodles available in the market. (Shin, 2003). Around the world, there are a plethora of

distinct instant noodle options (Seller et al. 2007). Indofood's Indomie instant noodles were the first instant noodle brand to be introduced in Nigeria in 1988. Numerous other nations throughout the world also distribute instant noodles (Seller et al. 2007). Nigeria has been producing instant noodles since 1995, however Indonesia is still the world's largest producer of indomie (World Instant Noodle Association (WINA, 2016).

Ever since the first instant noodle brand was introduced into Nigeria, rich people, middle-class people, and the impoverished have accepted it into many homes in both urban and rural areas. Children and even some adults referred to all types of instant noodles as "INDOMIE" as the indomie instant noodle developed a household word over time (Alabi et al. 2014). However, because of the intricate advertisements for many brands of these noodles, as well as their accessibility and ease of consumption, they have drawn people from a variety of social groups in both urban and rural areas, including children, adults, students, and the working class. 2010 saw Indofood release.

In routine clinical evaluations of the state of health, haematological measures are of diagnostic value and have been linked to health indices (Saliu et al. 2012). It is well known that long-term illnesses have a negative impact on blood cells. Blood parameter analysis is important for assessing the risk of changes to the human haematological system (Olson et al. 2000). Free radical damage causes oxidative DNA, protein denaturation, lipid peroxidation, disruption of membrane fluidity, and changes in platelet functions. These effects have been connected to a number of chronic health issues, including diabetes, cancer, inflammation, aging, and atherosclerosis. Despite the fact that nearly all organisms have evolved antioxidant defense and repair systems to shield them from free radicals, these systems are not enough to shield them entirely from oxidative damage (Nwanna and Oboh, 2007).

Glutamate consumption has been connected, independently of calorie intake and physical exercise, to obesity and metabolic syndrome. Additionally, several researchers have documented glutamate's neurotoxic consequences. Scientists generally agreed that glutamate is safe for use in children, expectant mothers, and nursing mothers. It has been demonstrated that, under typical situations, glutamate has a very low acute and chronic toxicity profile. More research is required to determine how this seasoning affects albino rats' organosomatic and haematological parameters, given the contradictions in the literature and the growing concerns about the safety of using monosodium glutamate.

MATERIALS AND METHODS

Indomie Seasoning

Indomie noodles (Chicken flavour) manufactured February 2023 by De United Foods industries Limited, Idiroko road, Ota, Ogun State Nigeria was bought at Nyong Essien Market Uyo Akwa Ibom State and the seasoning powder containing iodized salt, sodium polyphosphate, potassium bicarbonate, sodium bicarbonate, flavour enhancer (E621, E627, E631), Sugar, yeast extract, soy powder, ginger powder, garlic powder, chicken flavour was used for experiment.

Experimental Animals

The Laboratory Animal House, Department of Pharmacology, Faculty of Pharmacy, University

of Uyo, Nigeria, provided sixteen (16) normal, healthy female albino rats weighing between 83 and 140 g. They were housed in a clean, well-ventilated laboratory setting. The rats were given unlimited access to clean water and conventional rat food pellets after four weeks of acclimatization. The National Institute of Health's (1985) Guidelines for the Care and Use of Laboratory Animals and the University of Uyo's (Nigerian) Ethical Guidelines for the Use of Laboratory Animals were followed in all experimental methods.

Experimental Procedure

Following a month of acclimation, the rats were split into four groups, each consisting of four rats. As the control group, Group A was given normal saline (3 ml/kg). For 21 days, the seasoning powder dosages of 25 mg/kg, 50 mg/kg, and 75 mg/kg were administered to Groups B, C, and D, respectively. Every seasoning was taken orally. Diethyl ether was used to anesthetize the rats following their 21-day treatment. Every animal was killed by cervical dislocation, the heart was carefully accessed by cutting up the thoracic cage, and a cardiac puncture was used to get a blood sample. To avoid blood clotting, the blood from each rat was taken and placed into labelled 5ml heparinized Ethylene Diamine Tetra-acetic acid 8.5% (EDTA) vials.

Blood Sample: Rats were weighed and fasted for an entire night following their 21-day exposure before having blood samples taken. After that, the rats were killed. We collected all blood samples between 7 and 9 am in order to minimize variations resulting from circadian rhythm. Blood samples were quickly combined with the anticoagulant in EDTA (Ethylene Diamine Tetra-Acetic Acid 8.5%) anticoagulant tubes. After being labeled, every blood sample was taken right away to the lab for analysis.

Organosomatic index of Experimental rats

Throughout the experiments, weekly records of body weight changes were made. At the conclusion of the trial, the experimental rats' body weight gain and a few key organs were weighed. Using the Sellers et al. (2007) approach, the liver and kidney (left and right) were excised and weighed in order to determine the organosomatic index.

Organ - organ body wight ratio = $\frac{\text{Weight of the gain}}{\text{Weight of the whole animal}} \times 100$

Haematological parameters

The improved Nuebauerhemocytometer was used to count blood, and an automatic haematological assay analyzer was used to determine several haematological parameters, including packed cells (PCV), haemoglobin concentration (Hb), total red blood cell (RBC) count, mean cell volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), total white blood cell (WBC), and total platelets count. The following formula was used to determine the haematological indices, such as MCHC, MCH, and MCV, Dacie and Lewis (2001).

$$MCHC (fdl - 1) = \frac{Hb(gdl - 1)}{PCV(\%)} \times 100$$

$$MCH (pgcell-1) = \frac{Hb (gdl - 1)}{RBC \text{ count in millions } mm^{-3}} \times 100$$

$$MCV (flcell-1) = \frac{PCV\%}{RBC \text{ count in millions } mm^{-3}} \times 100$$

Haematological Analysis

Haematological parameters determination was carried out at the Haematology unit, Ibom


Specialist Hospital Uyo-Ikot Ekpene road, Ekit Itam Akwa Ibom State, Nigeria, using Automated Hematology Analyzer (Mindray BC-5380, UK –manufactured in 2017.

Statistical Analysis

Using the SPSS-PC software package (Version 26.0 SPSS Inc. Chicago, USA), the group means ±SEM for each parameter was computed, and significant differences were identified by Analysis of Variance (ANOVA). Means were separated using Turkey's Post hoc test at 5% level of significance. A clustered chart was utilized to evaluate the impacts on several days.

RESULTS

Figure 1 displays the body weights at the start of the trial (starting weight), at midpoint (2 weeks), and at final (3 weeks) following the oral administration of indomie seasoning to all experimental groups. When the experimental rats were given oral doses of the seasoning powder at 25 mg/kg, 50 mg/kg, and 75 mg/kg, their body weight increased significantly in comparison to the control group.



The liver's organ weight (hepatosomatic index) increased significantly ($p < 0.05$) in all groups when compared to the control, according to the findings in Table 1. Additionally, it demonstrates that there was no discernible difference in the organosomatic index in the liver and kidney between the control group and the


groups that received 25%, 50%, and 75% of their protein from Indomie noodles (groups A, B, and C). On the other hand, the organosomatic index differed significantly ($p < 0.05$) between the groups given indomie seasoning orally, and the weight of the right and left kidneys increased significantly ($p < 0.05$).

Table 1: Effects of indomie seasoning on some organosomatic index in female albino rats

Parameters	Mean±standard deviation			
	Group A (25%)	Group B (50%)	Group C (75%)	Control
Hepatosomatic index	0.80±0.00	0.80±0.00	0.80±0.10	0.84±0.20
Renosomatic index (Right)	4.20±0.40	3.81±0.30	4.24±0.40	4.14±0.50*
Renosomatic index (Left)	4.15±0.40	3.57±0.20	4.16±0.45	3.98±0.40*

Significance levels are indicated as follows: *indicates significance at the 5 % level

Clustered Bar Mean of Wbc, Mean of Neu, Mean of Lym, Mean of Mon, Mean of Eos, Mean of Bas by treatment by INDEX




The data presented in figure 2 demonstrated a significant difference ($p > 0.05$) between the WBC counts of albino rats given different amounts of indomie seasoning and the control groups.

The results, which are displayed in figure 3, demonstrated that the albino rats given Indomie Seasoning experienced a substantial shift in packed cell volume and a significant drop in mean

haemoglobin as compared to the control group. While not statistically significant, the mean corpuscular hemoglobin concentration dropped when compared to the control.


Clustered Bar Mean of Rbc, Mean of Hgb, Mean of HCT, Mean of MCV, Mean of MCH, Mean of MCHC, mean of RDW.CV, Mean of RDW.SD by treatment of INDEX



Keys: RBC – Red Blood Cells; HGB – Haemoglobin Concentration, HCT – Haematocrit Concentration, MCV – Mean Cell Volume, MCH – Mean Corpuscular Haemoglobin, MCHC – Mean Corpuscular Haemoglobin Concentration, RDW.CV – Red Cell Distribution Width RDW.SD- Red Cell Distribution Width

There was no discernible difference between the treated groups and the control when indomie seasoning was examined for its effect on albino rats' platelet levels. Groups 1 and 2 showed very little

differences between the treatment and the control, while group 2 demonstrated notable differences among the treated, as indicated in figure 4.



Keys: PLT – Platelets Concentration, MPV – Mean Platelet Volume, PDW – Platelet Distribution Width, PCT – Plateletcrit

DISCUSSIONS

The study's findings demonstrated that the experimental rats' body weights increased significantly across all groups when they were fed indomie seasoning orally. This is consistent with research by Hermanussen et al. (2006), who found that rats' body weight increased when they consumed monosodium glutamate, suggesting that the substance may be responsible for obesity and metabolic diseases. The results of this investigation are at odds with those of studies conducted by Mululy et al. (2013) and Tordoff et al. (2012), which found that consuming monosodium glutamate did not affect adult rats' or mice's body weight and was therefore unlikely to contribute to obesity.

When compared to the control group (group 4), the organosomatic index significantly changed

in all three treated groups (25 mg/kg, 50 mg/kg, and 74 mg/kg). According to Nelson et al. (2002), an increase in the organ-body weight ratio is a sign of inflammation. According to reports, an increase in an animal's liver weight is frequently indicative of induced toxicity brought on by the stimulation of liver enzymes; this toxicity may be brought on by a notable growth of the smooth endoplasmic reticulum (Anozie and Onwurah, 2001). Nephropathy may be indicated by the rise in kidney weight (Ijeh and Obidoa, 2001). Additionally, as observed in the cases of hyponatremia and hypernatremia, the cellular transporters activity is roughly altered by up or down regulates as brain tissue swells or shrinks. On the other hand, a variety of factors, such as acute hyponatremia, drug poisoning, water intoxication, and hypoxia, can result in either cell shrinkage or swelling. Brain enlargements have the potential to cause significant

cytotoxic oedema as well as a noticeable reduction in the size of the basal cisterns and ventricular system (Tordoff et al. 2012).

Animal blood is a biological fluid that takes metabolic waste products out of the cells and provides essential elements like nutrition and oxygen to the cells. Red blood cells, white blood cells, platelets, and plasma—the liquid component of blood that comprises water, proteins, salts, lipids, and glucose—are among the components of blood (Alberts, 2012). Blood is essential for sustaining homeostasis and controlling the body's functions. It performs a wide range of functions within the body, including the following: feeding tissues with oxygen (bound to haemoglobin, which is carried by red blood cells), supplying nutrients and fatty acids either dissolved in blood or bound to plasma proteins, eliminating waste, immune functions, including the circulation of white blood cells and the detection of foreign material by antibodies, coagulation, which is a component of the body's self-repair mechanism (blood clotting by the platelets after an open wound to stop bleeding), messenger functions, including the transport of hormones and the signalling of tissue damage, regulating body pH, regulating core body temperature, and hydraulic functions, among others (Martini and Nath, 2009).

Most haematological indicators showed significant variations between treatment groups. The addition of Indomie noodles in the meals did not cause any pathological effects, as indicated by the total leucocyte count, which suggests that the rats' health status was normal (Paul et al. 2012). In comparison to the control group, there was a substantial ($p < 0.05$) rise in the percentage of lymphocytes and a decrease in the percentage of monocytes and granulocytes. Examining the red blood cell indices revealed lower MCV, MCH, and MCHC levels. This demonstrates that all blood parameters showed differences between the three levels of Indomie noodles (25, 50, and 75%) and the control values. The reduced ability of the bone marrow to produce haemoglobin is most likely the cause of the corpuscles' decreased haemoglobin levels (Sharma, 2015). When compared to the control group, a 75% rise in platelet count may indicate secondary thrombocytosis. It was proposed that this rise might be the result of consuming excessive amounts of salt and glutamate, two ingredients in Indomie seasoning

that have an impact on the body's water balance and blood and bodily fluid compartments. They go on to say that it might have something to do with the thrombopoietin hormone's biosynthesis, which increases platelet production (Tordoff et al. 2012).

The evaluation of haematological parameters can be employed to ascertain the degree of detrimental impact that foreign substances, such as plant extract, have on activities associated to blood. In comparison to their respective control groups, Yakubu et al. (2007) found that a prolonged administration of the alcoholic bitters significantly reduced the levels of haemoglobin, red blood cells, packed cell volume, platelets, and white blood cells, among other differential counts.

According to the current study, the neutrophil count decreased significantly at all seasoning doses, and the decrease was greater in the groups who got the highest dose. Human neutrophil counts typically range from 3000 to 6000 / cu.mm of blood.

This could mean that the seasoning has a time-dependent, direct toxic effect on blood neutrophils or that it negatively affects bone marrow blood formation, particularly on progenitor cells (aplasia). The initial line of defense against foreign chemicals, poisonous substances, and invasive microorganisms is provided by neutrophils and monocytes, highlighting the critical function neutrophils play in the body's defense (Hall, 2011). This could be a sign of the treated rat groups' immunological condition declining due to the seasoning's harmful effects. All treated animals showed an observed increase in lymphocyte count (normal value = 1500 to 2700 / cu.mm of blood). This increase could be attributed to the animals' perception of the seasoning as a toxic agent, a significant increase in granulocytes, or the interaction between MSG and gastrointestinal macrophages. Macrophages function as antigen-presenting cells, delivering antigenic components (polypeptides) to activate B cells and helper T cells (Sembulingham, 2005). Additionally, macrophages emit chemicals known as interleukin-1 / -cytokines, which cause the lymphocyte count to rise and the lymphocytes to proliferate (Sembulingham, 2005; Barrett et al. 2010). MSG did not have a deleterious impact on lymphocytic cells, as evidenced by the steady increase in counts observed in all withdrawal groups as well as in Groups B1 and B2, which

received therapy for extended periods of time. This suggests that MSG has an ongoing, lingering impact on the body's systems.

The seasoning had a decreasing effect on both the concentration of hemoglobin and the packed cell volume. Humans typically have packed cell volume of 38–42%, hemoglobin concentration of 14–16 g%, and red blood cell count of 4–5.5 million/cu.mm, respectively. RBC count indicates that it most likely shortens red blood cell life in the blood, maybe due to direct toxicity. This could have potentially been mediated by negatively affecting the bone marrow's hematopoietic stem cells. Increased oxidative stress, a byproduct of anaerobic respiration, may be brought on by MSG in the animal tissues. Micronucleated polychromatic erythrocytes (MNPCEs) were considerably ($p < 0.01$) stimulated to develop by 4 mg/g of MSG (Farombi and Onyema, 2006). Elphick et al. (2008) used HT22 murine hippocampal cells as a model for oxidative stress-induced neuronal death to examine the oxidative glutamate toxicity. According to Elphick et al. (2008), these harmful effects were typified by nuclear and cell condensation but did not result in DNA fragmentation or the release of cytochrome c from the mitochondria. Significant increases were seen in the MCV and MCH levels (normal values = 27 to 32 pg and 78 to 90 cu. micron, respectively). Although there was no discernible ameliorating effect, the elevated values are more suggestive of anemia.

Macrocytic cells are indicated by an increased mean corpuscular volume (MCV) (Sembulingham, 2005). While elevated MCH is suggestive of macrocytic anaemia, high MCV is reported in pernicious anaemia (normochromic) and megaloblastic anaemia (hypochromic) (Richards, 1993). Nonetheless, both the treatment and control groups have normal mean corpuscular haemoglobin concentrations (MCHCs; typical values for humans are between 30 and 38%). More precisely, macrocytic normochromic anaemia, or pernicious anaemia, was recommended. This could be because of the acidic MSG (L-form of glutamic acid) that causes the gastric mucosa to atrophy (gastritis). This leads to a decrease in the production of the intrinsic factor and poor absorption of vitamin B12, which is the primary cause of pernicious anaemia (Sembulingham, 2005).

The current study demonstrates that the

administration of indomie seasoning significantly affects the neutrophil and lymphocyte counts, which are suggestive of poisoning and weakened immune systems, respectively, in the treated animals. While changes in the treated animals' PCV, Hb, RBC, MCV, and MCH values were all suggestive of anaemia. Therefore, our results agree with Eweka, (2007) who stated that monosodium glutamate is unhealthy even when it serves as a flavouring agent.

CONCLUSION

Due to its association with Chinese Restaurant Syndrome (CRS), this study showed that indomie noodle seasoning containing monosodium glutamate may be harmful to human health. Long-term, frequent use of seasonings containing monosodium glutamate can cause diseases like obesity, kidney impairment, fibroids, and hepatotoxicity. To educate people about the dangerous effects of the indomie noodle seasoning, more awareness should be raised, and natural substitutes for the seasoning should be encouraged.

REFERENCES

- Alabi AO, Edward BE, Oluwatobi CO, Olutayo SS. (2014). Assessment of the mutagenic and genotoxic potential of indomie noodle seasoning using bacterial (Salmonella) reverse mutation and chromo tests. *Journal of Innovative Biology*. (1): 210-214.
- Alberts B. (2012). *Molecular Biology of the Cell*. NCBI Bookshelf. ISBN-10-0-8153-4072-9.
- Amin KA, Abdel-hameid H, Abd-elsttar AH. (2010). Effect of food azo dyes tartrazine and carmoisine on biochemical parameters related to renal, hepatic function and oxidative stress biomarkers in young male rats. *Journal of Food and Chemical Toxicology*. 36(48): 2994-2999.
- Anozie OI, Onwurah IN. (2001). Toxic effects of Bonny light crude oil in rats after ingestion of contaminated diet. *Nigerian Journal of Biochemistry and Molecular Biology*. 16(3):1035-1085.
- Barrett KE, Barman SM, Boitano S, Brooks HL. (2010). *Ganong's Review of Medical Physiology*. 23rd Edition, pp. 63-78.
- Dacie JV, Lewis SM. (2001). *Practical Haematology*. 11th Edn., Longman Group Ltd., Hong Kong. pp: 11-17.
- Elphick LM, Hawat M, Toms NJ, Meinander A, Mikhailov A, Eriksson JE. (2008). Department of Biochemistry and

- Pharmacy, Abo Akademi University, FIN-20521 Turku; Kass, George E. 2008-01-01.
- Eweka AO. (2007). Histological studies of the effects of monosodium glutamate on the kidney of adult Wistar rats. *Internet Journal of Health*. **6(2)**. (www.ispub.com).
- Farombi EO, Onyema OO. (2006). Monosodium glutamate-induced oxidative damage and genotoxicity in the rat: modulatory role of Vitamin C, Vitamin E and quercetin. *Human Experimental Toxicology*. **25(5)**: 251-259.
- Hall JE. (2011). Guyton and Hall Textbook of Medical Physiology. 12th Edition, Philadelphia, pp. 423-431.
- Hermanussen M, Garcia AP, Sunder M, Voigt M, Salazar V, Tresguerres JAF. (2006). Obesity, voracity and short stature: the impact of glutamate on the regulation of appetite. *European Journal of Clinical Nutrition*. **60**: 25-31.
- Hope N. (2001). CNN.com. Instant noodles a health hazard. A report. Retrieved 7th November, 2012.
- Ijeh II, Obidoa O. (2001). Effects of dietary incorporation of two varieties of *Vernonia amygdalina* Del leaves on relative organ weights of weaning albino rats. *Nigerian Journal of Biochemistry and Molecular Biology*. **6**.
- Indofood (2010). Total Food Solutions Company Limited.
- Josiah IE, Clark PE, Lin IC. (2017a) Utilization of noodle waste as replacement for maize in the diet of broiler starter chickens. Cooking time of white salted noodles and its relationship with protein and amylose contents of wheat. *Cereal Chemistry*. **81**: 165-171.
- Kim SG, Instant noodles. (1996a). In: Pasta and Noodle Technology; Kruger J.E., Matsuo, R.B, Dick J. W, (Eds.) American Association of Cereal Chemistry, St Paul, MN, USA. pp. 195-225.
- Maluly HDB, Areas, M. A., Borelli, P. and Reyes, F. G. R. (2013). Evaluation of biochemical, hematological and histological parameters in non-diabetic wistar rats fed monosodium glutamate. *Food and Nutrition Sciences*. **4**: 66-76.
- Martini F, Nath JI. (2009). Fundamentals of Anatomy and Physiology. Nath Judi Lindsley (8th edition). San Francisco: Pearson/Benjamin Cummings, 657p.
- National Institute of Health (NIH). Guide for the use of laboratory animals. DHHS, PHS, NIH Publication No. 85-23 (1985 Revised).
- Nelson G, Chandrashekar J, Hoon MA, Feng L, Zhao G. (2002). An amino acid taste receptor. *Nature*. **416**: 199-202.
- Nwanna EE, Oboh G. (2007). Antioxidant and hepatoprotective properties of polyphenol extracts from *Telfairia occidentalis* (Fluted pumpkin) leaves on acetaminophen induced liver damage. *Pakistan Journal of Biological Sciences*. **10(16)**: 2682-2687.
- Olson H, Betton G, Robinson D, Thomas K, Monro A, Kolaja G, Lilly P, Sanders J, Sipes G, Bracken W, Dorato M, Van Deun K, Smith P, Berger B, Heller A. (2000). Concordance of the toxicity of pharmaceuticals in humans and in animals. *Toxicology and Pharmacology*. **32**: 56-67.
- Paul MV, Abhilash M, Varghese MV, Alex M, Nair RH. (2012). Protective effects of alpha-tocopherol against oxidative stress related to nephrotoxicity by monosodium glutamate in rats. *Toxicological and Mechanical Methods*. **22(8)**: 625-630.
- Richards GE. (1993). Winthrobe's Clinical Hematology. 9th edition, 1: 715-745.
- Saliu JA, Elekofehinti OO, Komolafe K, Oboh G. (2012). Effects of some green leafy vegetables on the haematological parameters of diabetic rats. *Journal of Natural Products and Plant Resources*. **2(4)**: 482-485.
- Sellers RS, Morton D, Michael B, Roome N, Johnson JK, Yano BI, Perry R, Schafer K. (2007). Society of Toxicological Pathology position paper: Organ-weight recommendations for toxicology studies. *Toxicological Pathology*. **35**: 751-755.
- Sembulingham K. (2005). Essentials of Medical Physiology. 2nd edition pp. 46-56.
- Sharma A. (2015). Monosodium glutamate-induced oxidative kidney damage and possible mechanisms: a mini-review. *Journal of Biomedical Science*. **22**: 93.
- Shin SN, Kim SK. (2003). Properties of instant noodle flours produced in Korea. *Cereal Foods World*. **48(6)**: 310-314.
- Tordoff MG, Aleman TR, Murphy MC. (2012). Effects of monosodium glutamate consumption on the body weight or composition of adult rats and mice. *Journal of Physiology of Behaviour*. **107**: 338-345.
- World Instant Noodle Association (WINA) (2016). Expanding Market: Global demand for instant noodles. <https://instantnoodleres.org> Retrieved 8th February.
- Yakubu MT, Akanji MA, Oladiji AT. (2007). Haematological Evaluation in male albino rats following chronic administration of aqueous extract of *Fadogia agrestis* stem. *Pharmacognosy Magazine*. **3(9)**: 34-38.