



PHYSICOCHEMICAL PROPERTIES AND MICROBIAL STABILITY OF COMPOSITE FLOUR PRODUCED FROM RICE AND UNRIPE PLANTAIN FOR *TUWO* PRODUCTION

Mutiat A. B.^{1*}, Fausat L. K.¹, Sarafa A. A.^{2,3*}, Titilayo N. A.¹, Itohan S.A.^{2,4}

¹Department of Home Economics and Food Science, Faculty of Agriculture, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria

²Department of Agricultural Technology, School of Agricultural Technology and Engineering, Edo State College of Agriculture and Natural Resources, Iguoriakhi, Edo State, Nigeria

³Department of Food Technology, Faculty of Technology, University of Ibadan, Ibadan, Oyo State, Nigeria

⁴Department of Microbiology, Faculty of Life Sciences, University of Benin, PMB 1154, Benin City, Nigeria

*Author for Correspondence: akeemsarafa@yahoo.com

ABSTRACT

Tuwo shinkafa, a dumpling widely consumed in the northern part of Nigeria, is usually produced from non-parboiled, soft, mashed white rice. With the aim of enhancing its nutritional composition and making it suitable for consumers suffering from diabetes, obesity and other related health issues, the suitability of rice and unripe plantain composite flour for *tuwo* production was examined. Rice and unripe plantain flour were produced and mixed at varying ratios; 100:0 (control), 95:5, 90:10, 85:15 and 80:20. The proximate, functional properties and microbial stability of the composite flour and sensory acceptability of the *tuwo* produced were investigated. The ash (0.88-3.76%), fibre (5.63-6.06%), swelling capacity (218.30-236.90%) and water absorption capacity (178.40-194.80%) of the rice-unripe plantain composite flour increased while the packed bulk density (0.18-0.32 g/ml) decreased with increase in the level of plantain flour substitution. The 100% rice flour exhibited the highest ($p < 0.05$) oil absorption capacity (65.83%). Although no bacterial growth was observed on all the fresh flour, the bacterial and fungal counts of the flour increased with increase in the level of plantain flour substitution and also as the storage duration increased. This study revealed that up to 20% unripe plantain flour could be substituted in rice flour for the production of highly nutritive, wholesome and acceptable *tuwo shinkafa*, though 10% substitution is recommended based on the sensory ratings.

Keywords: Unripe plantain flour, food safety, microbial stability, physicochemical properties, acceptability, rice flour, *tuwo shinkafa*

INTRODUCTION

Tuwo shinkafa is a popular meal in the northern part of Nigeria from where it is believed to have originated. It is usually made by mixing non-parboiled, soft, sticky white rice flour with boiling water and stirred until a smooth consistency gel is formed. This gel, moulded into a spherical shape, is usually consumed with accompaniment (Falade and Christopher, 2015) including soups such as *miyan kuka*. Rice (*Oryza sativa*) is a cereal of great importance in both developing and developed worlds. It is a semi-aquatic, annual grass plant which has been reported to contain substantial amount of carbohydrate, vitamins such as thiamine, niacin and riboflavin, minerals and little amount of protein and fat (Fresco, 2005; Umadevi *et al.*, 2012). It is also gluten and cholesterol free. Oko *et*

al. (2012) showed that rice contained substantial amount of calcium, magnesium and phosphorus with little quantities of iron, copper, zinc and manganese. Rice is being regarded as the queen among cereals owing to its relatively good nutritional quality and digestibility (Anjum *et al.*, 2007). Freshly harvested rice grains contain about 80% carbohydrate which is made up of starch, glucose, sucrose and dextrin (Verma and Srivastav, 2017).

Rice is grown in all the ecological zones of Nigeria, with different varieties possessing adaptation traits suited to each ecological zone (Sanni *et al.*, 2005). In Nigeria, *Oryza sativa* (Asian rice) and *Oryza glaberrima* (African rice) are the two commonly cultivated varieties of rice out of which *O. sativa* is the most widely cultivated (Abulude, 2004; Adeyemi *et al.*, 1986). *Ofada* rice

is the most cultivated rice in Nigeria especially in the South-western agro ecological zone and annual production of rice in Nigeria was estimated at about 3 million tonnes (Adebowale *et al.*, 2010). The production and consumption of rice as a major dietary source of energy is at large in Nigeria and the use of rice flour as a staple food is on the increase worldwide. This staple food provides 700 calories/day-person for about 3000 million people of the world's population (Vlachos and Arvanitoyannis, 2008). Despite the nutritional value and worldwide consumption of rice and its products, its relatively high glycemic index which poses major threats to diabetics and dieters, is an impediment to its utilization.

Plantain (*Musa paradisiaca AAB*) is a popular dietary monocotyledonous perennial and most important staple crop in the tropical and sub-tropical regions of the world after rice, wheat and maize (Baiyeri *et al.*, 2011; Kawongolo, 2013). It is a highly perishable climacteric fruit which when harvested at the mature but unripe stage ripens within two to seven days (Abiodun-Solanke and Falade, 2011). It is a source of essential nutrients and income for many households around the world including those in sub-Sahara Africa (Kawongolo, 2013) and Africa contributes over 12 million metric tons to the over 12 million metric tons annual world production of plantain. Nigeria is the biggest producer of plantains in West Africa with an estimated production of about 2,722,000 metric tons in 2009, majority of which were produced and harvested from the southern part of the country, with an average consumption level of 190 kg/person/year (FAO, 2009; 2011).

Unripe plantain is well recognised for its richness in dietary fibre, resistant starch, vitamins and minerals with low quantities of protein and fat (Ayodele and Erema, 2010; Baiyeri *et al.*, 2011; Agu and Okoli, 2014; Karim *et al.*, 2020). Unripe plantain products have been reported to exhibit low glycemic index and blood glucose response (Ayodele and Erema, 2010). This indicates that unripe plantain diets would be ideal for diabetics, dieters and pregnant women. The dietary management of diabetes, obesity and other related health challenges could therefore be achieved through consumption of diets containing plantain flour. The inclusion of plantain flour in food formulation will not only improve the nutrients intake of the consumers but will also enhance the

utilization of plantain and create varieties in human diet. Previously, a study has reported the production of acceptable dumpling dough from *ofada* rice and unripe plantain flour with appropriate dietary fibre ratio, cholesterol reduction and low weight gain potentials (Arueya and Akande, 2018) but existing gap relating to the keeping quality of the developed flour blends and acceptability of the final product still exists in the literature. Consequently, this study was designed to determine the effect of partial substitution of unripe plantain flour on the physicochemical properties and microbial stability of rice flour as well as to evaluate the sensory acceptability of cooked paste or dumpling (*tuwo shinkafa*) produced from the composite flour.

MATERIALS AND METHODS

Materials

Local rice (*ofada funfun / Oryza sativa L.*) and fresh matured unripe plantains (*Agbagba*) at stage 1 of ripening (Aurore *et al.*, 2009) were purchased at Oja-oba market in Ilorin and carefully transported to laboratory of the Department of Home Economics and Food science, University of Ilorin, Nigeria for processing. All chemicals and the equipment used were of analytical grade.

Rice flour production

The method of Ogunlakin *et al.*, (2014) was adopted with little modification. The local rice was sorted, washed and cleaned. The cleaned rice was conditioned for 5 minutes and oven dried at 60°C for 4 hours. The dried rice was milled, cooled, sieved through 100 mesh sieve and packaged in polyethylene bag.

Unripe plantain flour production

Plantain flour was produced according to Yarkwan and Uvir (2015) with little modification. Fingers were picked and washed in a bowl of portable water to remove the latex and avoid darkening of the pulp during peeling. The washed fingers were peeled and then sliced in thin slices of about 5.0 mm thick using a stainless steel knife. The slices were oven-dried at 60°C for 24 hours after which they were cooled, milled, sieved into fine flour using a mesh net of about 0.2 mm mesh size and then packaged in polyethylene bag.

Formulation of composite flour

Composite flour for *tuwo* production was formulated from rice and unripe plantain flour as shown in Table 1.

Table 1: Formulation of rice and unripe plantain composite flour

Sample	Rice flour (%)	Unripe plantain flour (%)
R ₁₀₀ P ₀	100	0
R ₉₅ P ₅	95	5
R ₉₀ P ₁₀	90	10
R ₈₅ P ₁₅	85	15
R ₈₀ P ₂₀	80	20

Proximate composition of rice-unripe plantain composite flour

The moisture, dry matter, ash and crude fibre contents of the flour samples were evaluated using standard methods (AOAC, 2005). Briefly, oven drying method at 105 °C for 5 hours for moisture evaluation, dry matter was obtained by subtracting the percentage moisture content from hundred, total ash determination by igniting 2 g of each sample at 550 °C for 4 hours using muffle furnace and crude fibre was determined using digestion method.

Functional properties of rice-unripe plantain composite flour

Packed bulk density

Five grams (5 g) of each sample was weighed into a 25 mL graduated measuring cylinder. The cylinder was gently tapped on the laboratory table several times until there was no further diminution of the sample level. The volume of the sample was taken and the tapped bulk density was calculated thus (Onwuka, 2005);

$$\text{Packed bulk density (g/ml)} = \frac{\text{Weight of sample}}{\text{volume of sample after tapping}}$$

Swelling capacity

The method described by Ukpabi and Ndimele (1990) was used with modifications for the determination of swelling capacity of the rice-plantain composite flour. Five grams of flour sample was transferred into a 50 mL graduated

measuring cylinder. The sample was gently levelled by tapping the cylinder and the initial volume was recorded. Fifty millilitre of distilled water was poured into the cylinder. The cylinder was covered, inverted for proper mixing and then allowed to stand for 4 hours. The swelling capacity was taken as the percent multiple of the initial volume.

Water and oil absorption capacity

The method of Sosulski *et al.* (1976) was adopted in determining the water and oil absorption capacity of the flour. One gram of flour sample was mixed with 10 mL distilled water or 10 mL refined soybean oil (specific gravity, 0.9092) in a 50 mL centrifuge tube for water and oil absorption capacity, respectively. The mixture was allowed to stand at room temperature for 30 min and then centrifuged at 2000 ×g for 30 min. The percentage water or oil bound per gram flour was recorded as the water and oil absorption capacity, respectively.

Storage and microbial analysis of rice-unripe plantain composite flour

Microbial analysis was carried out on each flour sample. Then, a portion of each flour samples was transferred into different polyethylene bags and stored at room temperature (27±2°C) for 8 weeks during which microbial analysis was being carried out fortnightly.

The total viable bacterial and fungal counts of the flour were investigated using pour plate method (Aruwa and Akinyosoye, 2015). One gram of each flour sample was aseptically weighed and mixed with 9 mL of sterilized distilled water in a test tube. One millilitre of each suspension was taken for serial dilution (10⁻² and 10⁻⁴) and 1 mL aliquots were inoculated into sterile Petri dishes containing nutrient agar and potato dextrose agar, which have been prepared according to the manufacturers' instructions, for the enumeration of total bacteria and fungi, respectively. The nutrient agar's plates were incubated at 37°C for 24 hours while potato dextrose agar plates were incubated at room temperature (26 ± 2°C) for 3–5 days. The colonies were then counted using the Stuart scientific colony counter and expressed as colony forming units per gram (cfu/g) of flour.

Preparation and sensory acceptability of *tuwo shinkafa* produced from rice-unripe plantain composite flour

The method described by Bolade *et al.* (2002) for production of maize *tuwo* was adopted for the preparation of rice-plantain *tuwo shinkafa*. The overall ratio of flour to water used was 1:3.5 (w/v). Twenty five percent (25%) of the water was initially mixed with 20% of the measured flour to form slurry. Sixty percent of the measured water was heated to boiling point and the initially-prepared flour slurry was gradually added to the boiling water with continuous stirring until a gel-like consistency was obtained. The remaining flour (80%) was then added gradually to the boiling gel-like mass, with continuous stirring, until a satisfactory gel was obtained. The last quantity of water (15%) was added to the gel and covered for about 5 min without stirring (for effective cooking). It was vigorously stirred again and ready to be served. The resulting rice-based final product obtained is termed *tuwo shinkafa*.

The sensory qualities of *tuwo shinkafa* produced from rice and plantain composite flour were evaluated using multiple comparison test (Akeem *et al.*, 2023). The acceptability of the products in terms of colour, taste, mouldability, aroma, texture and overall acceptability was assessed by 30 untrained panellists comprising of students and members of staff of University of Ilorin, based on a 9-point hedonic preference scale (1 = dislike extremely, 2 = dislike very much, 3 = dislike moderately, 4 = dislike slightly, 5 = neither like nor dislike, 6 = like slightly, 7 = like moderately, 8 = like very much and 9 = like extremely). Drinkable water was provided for the panellists to rinse their mouths after evaluating each randomly presented coded sample.

Statistical analysis

All experiments were conducted in triplicates except where it is stated otherwise. The data were subjected to one-way analysis of variance (ANOVA) and significant difference among means was determined by Duncan's multiple range test ($p < 0.05$) using SPSS software version 15.0 (SPSS Inc., Chicago, IL).

RESULTS AND DISCUSSION

Proximate composition of rice-unripe plantain composite flour

The influence of partial substitution of rice flour with unripe plantain flour on the moisture, dry matter, ash and fibre contents is shown in Table 2. Moisture content is an important quality attribute on which the shelf stability and microbial growth susceptibility of any food depend. The moisture content of the flour blends ranged from 7.59-8.79% for 5% and 10% unripe plantain flour substituted samples, respectively. Although significant ($p < 0.05$) difference existed between 100% rice flour and the unripe plantain flour substituted samples, the moisture contents of all the flour samples were below 10%. The low moisture content obtained for the flour samples in this study agreed with the recommended standard of SON (2007) which stated that long term safe keeping of flour require the moisture level to be below 10%. This implied that the flour samples could be stored for a long time without biochemical or microbial deterioration. The solid component of the flour samples ranged between 91.21% and 92.41% for 10% and 5% unripe plantain flour substituted samples, respectively. Substitution of unripe plantain flour (5-20%) in rice flour significantly ($p < 0.05$) affected the dry matter content of the flour blends.

The ash content recorded for 100% rice flour was 0.88% and this was observed to increase significantly ($p < 0.05$) with increase in the level of unripe plantain flour substitution. The ash content obtained for 100% rice flour was similar to 0.80% reported by Juliano and Bechtel (1985), mean value of 0.99% reported for 20 rice varieties (Oko *et al.*, 2012) and fell within the range of 0.39-0.90% reported for six rice varieties marketed in Penang Island, Malaysia (Thomas *et al.*, 2013). The increment in ash content of unripe plantain flour substituted samples could therefore be attributed to high ash content of plantain flour. This is plausible since ash content as high as 5.44% has been reported for oven-dried unripe plantain flour (Yarkwan and Uvir, 2015). Ash content is a nutritional component that reflects the total amount of inorganic matter or mineral composition of food. Previous studies had shown that plantain contained high amount of essential minerals such as potassium, calcium, phosphorus and iron, and

various vitamins such as A, B₁, B₂ and C (Chandler, 1995; Karim *et al.*, 2020). The inclusion of unripe plantain flour in rice flour could therefore be employed as a strategy to tackle micronutrients deficiency in both developing and developed countries.

The crude fibre content (5.63-6.06%) of the flour samples generally increased with increase in the level of unripe plantain flour substitution, though significant ($p > 0.05$) differences were only recorded after 10%, 15% and 20% substitution

levels. This increment could be due to the high fibre content (10.11%) of the oven-dried unripe plantain flour as reported by Yarkwan and Uvir (2015). Dietary fibre is a functional nutritional component of great importance in human diet due to its ability to increase faecal output, lower risk of bowels disorders, faecal pH, blood pressure and reduce incidence of colon cancer, coronary heart disease, stroke, diabetes, obesity, blood pressure as well as certain gastrointestinal diseases (Anderson *et al.*, 2009).

Table 2: Proximate composition of rice-unripe plantain composite flour

Sample	Moisture (%)	Dry matter (%)	Ash (%)	Fibre (%)
R10P0	7.91 ^c	92.09 ^b	0.88 ^e	5.63 ^d
R9P5	7.59 ^d	92.41 ^a	1.64 ^d	5.64 ^d
R9P10	8.79 ^a	91.21 ^d	2.14 ^c	5.72 ^c
R8P15	8.03 ^b	91.97 ^c	3.20 ^b	5.86 ^b
R8P20	7.60 ^d	92.40 ^a	3.76 ^a	6.06 ^a

Values are means of triplicate determinations. Means with the same superscript along the same column are not significantly ($p > 0.05$) different.

R₁₀₀P₀ = 100% rice flour; R₉₅P₅ = 95% rice flour + 5% unripe plantain flour; R₉₀P₁₀ = 90% rice flour + 10% unripe plantain flour; R₈₅P₁₅ = 85% rice flour + 15% unripe plantain flour; R₈₀P₂₀ = 80% rice flour + 20% unripe plantain flour

Functional properties of rice-unripe plantain composite flour

The packed bulk density (0.18-0.32 g/ml), swelling capacity (218.30-236.90%), water (178.40-194.80%) and oil (62.84-65.83%) absorption capacities of rice-unripe plantain composite flour are presented in Table 3. Substitution of unripe plantain flour (5-20%) in rice flour resulted in significant ($p < 0.05$) decrease in packed bulk density of the flour blends. Similar trends have been reported for the bulk density of wheat-plantain flour (Ogunlakin *et al.*, 2014), wholemeal wheat-unripe plantain flour (Inyang and Asuquo, 2016), plantain-tigernut flour (Adegunwa *et al.*, 2017) and *ofada* rice-plantain flour (Arueya and Akande, 2018), indicating higher particle compactness of plantain flour. Bulk density of a food material is essential in determining material handling, packaging requirement and food

applications (Adebowale *et al.*, 2008) since it depends on individual particle's mass, size, property, density and geometry (Kolawole *et al.*, 2016). The lower bulk density of unripe plantain flour substituted samples could be advantageous in the formulation of complementary foods as postulated by Akpata and Akubor (1999).

The swelling capacity and water absorption capacity of the composite flour generally increased with increase in the level of unripe plantain flour substitution. This was similar to the report of Inyang and Asuquo (2016) for wholemeal wheat and unripe plantain composite flour. Arueya and Akande (2018) also observed increase in swelling capacity with increase in the levels of plantain substitution in *ofada* rice. Swelling capacity is associated with binding within the starch granules of the micelle network (Kolawole *et al.*, 2016) and the increment in swelling capacity of the unripe

plantain flour substituted samples could be due to lower amylose and higher amylopectin content of plantain flour compared to rice flour.

Water absorption capacity reflects the ability of a food material to associate with water under limited water condition with the aim of improving handling (Giami and Bekebian, 1992). Similarly, increase in water absorption capacity with increase in plantain flour substitution in wholemeal wheat flour (Inyang and Asuquo, 2016) and tigernut flour (Adegunwa *et al.*, 2017) have been reported by previous researchers. The increment in water absorption capacity of unripe plantain flour substituted samples might be associated with presence of hydrophilic amino acids, dietary fibre and low amylose to amylopectin ratio in the unripe plantain flour. The higher water absorption capacity obtained for the rice-unripe plantain composite flours suggested that they would be very useful for aqueous food formulation such as bakery products which need proper hydration.

The oil absorption capacity of the composite flour decreased disproportionately with increase in the level of substitution of unripe plantain flour. The decrease in oil absorption capacity of the flour blends following incorporation of plantain flour was similar to the observation of Inyang and Asuquo (2016) on wholemeal wheat-plantain composite flour. The oil binding capacity of a food material is an indicator of its ability to absorb and retain oil, which consequently affects the flavour, mouthfeel, palatability and shelf life of food products such as bakery, soup mixes or meat products where fat absorptions are desired (Seena and Sridhar, 2005). The differences in oil binding capacity could be as a result of variations in the presence of non-polar side chains, which might bind the hydrocarbon side chains of oil among the flours (Adebowale and lawal, 2004).

Table 3. Functional properties of rice-unripe plantain composite flour

Sample	Packed bulk density (g/ml)	Swelling capacity (%)	Water absorption capacity (%)	Oil absorption capacity (%)
R ₁₀₀ P ₀	0.32±0.00 ^a	218.30±0.35 ^c	178.40±0.27 ^c	65.83±1.32 ^a
R ₉₅ P ₅	0.26±0.00 ^b	219.70±0.60 ^{bc}	184.80±2.91 ^b	62.84±2.0 ^{ab}
R ₉₀ P ₁₀	0.22±0.00 ^c	230.10±3.36 ^{abc}	189.20±4.06 ^{ab}	65.19±1.18 ^{ab}
R ₈₅ P ₁₅	0.21±0.00 ^d	233.50±2.37 ^{ab}	189.30±1.11 ^{ab}	63.25±0.76 ^{ab}
R ₈₀ P ₂₀	0.18±0.00 ^c	236.90±11.50 ^a	194.80±0.30 ^a	63.93±2.27 ^{ab}

Values are means of triplicate determinations ± SD. Means with the same superscript along the same column are not significantly ($p > 0.05$) different.

R₁₀₀P₀ = 100% rice flour; R₉₅P₅ = 95% rice flour + 5% unripe plantain flour; R₉₀P₁₀ = 90% rice flour + 10% unripe plantain flour; R₈₅P₁₅ = 85% rice flour + 15% unripe plantain flour; R₈₀P₂₀ = 80% rice flour + 20% unripe plantain flour

Microbial stability of rice-unripe plantain composite flour

Food safety, acceptability, shelf stability and fitness for consumption depend largely on its microbial stability. The total bacterial and fungal counts of rice-unripe plantain composite flour during 8 weeks storage are shown in Tables 4 and 5, respectively. No bacterial growth was observed on

the fresh flour samples. After two weeks of storage, bacterial growth ($1.0 - 2.30 \times 10^4$ cfu/g) was recorded for the flour samples and this growth was observed to increase with increase in the level of unripe plantain flour substitution and storage time up to the 8 weeks storage period ($2.20 - 3.50 \times 10^4$ cfu/g). The result was similar to that of Oviasogie *et al.* (2016) who reported that the total bacterial

count (cfu/g) of all wheat-plantain flour increased with increasing level of plantain flour during the nine weeks period of storage. High bacteria counts could be an indication of potential health hazards and food spoilage.

The fungal count of the rice-unripe plantain flour samples varied between 2.00×10^4 cfu/g and 4.60×10^4 cfu/g within the 8 weeks of storage. Similar to bacterial count, the fungal count of the composite flour increased with increase in the level of unripe plantain flour substitution and storage time. This result was also similar to that of Oviasogie *et al.* (2016) who reported increase in the fungal growth with increasing levels of plantain

flour substitution and storage time. These implied that substitution of unripe plantain flour in rice flour enhance the susceptibility of the resulting composite flour to microbial contamination and growth. The rapid microbial proliferation on the flour samples could be attributed to nutrient availability and favourable micro-environment with resultant recovery of homeostatic imbalance (Ogbulie *et al.*, 1993). Fungal growth has been associated with the formation of heat stable mycotoxins which is a major concern for food quality and safety (Dalié *et al.*, 2010, Tang *et al.*, 2019).

Table 4: Total bacterial count of rice-unripe plantain composite flour during 8 weeks storage period

Sample	Week 0 ($\times 10^4$ cfu/g)	Week 2 ($\times 10^4$ cfu/g)	Week 4 ($\times 10^4$ cfu/g)	Week 6 ($\times 10^4$ cfu/g)	Week 8 ($\times 10^4$ cfu/g)
R100P0	-	1.0	1.6	2.0	2.2
R95P5	-	1.4	1.8	2.3	2.6
R90P10	-	1.8	2.0	2.5	2.9
R85P15	-	2.0	2.2	2.7	3.1
R80P20	-	2.3	2.6	3.0	3.5

R₁₀₀P₀ = 100% rice flour; R₉₅P₅ = 95% rice flour + 5% unripe plantain flour; R₉₀P₁₀ = 90% rice flour + 10% unripe plantain flour; R₈₅P₁₅ = 85% rice flour + 15% unripe plantain flour; R₈₀P₂₀ = 80% rice flour + 20% unripe plantain flour

Table 5: Total fungal count of rice-unripe plantain composite flour during 8 weeks storage period

Sample	Week 0 ($\times 10^4$ cfu/g)	Week 2 ($\times 10^4$ cfu/g)	Week 4 ($\times 10^4$ cfu/g)	Week 6 ($\times 10^4$ cfu/g)	Week 8 ($\times 10^4$ cfu/g)
R ₁₀₀ P ₀	2.0	2.2	2.6	2.8	3.4
R ₉₅ P ₅	2.3	2.5	2.9	3.2	3.8
R ₉₀ P ₁₀	2.5	2.7	3.2	3.6	4.0
R ₈₅ P ₁₅	2.7	3.0	3.5	3.9	4.2
R ₈₀ P ₂₀	3.0	3.3	3.6	4.2	4.6

R₁₀₀P₀ = 100% rice flour; R₉₅P₅ = 95% rice flour + 5% unripe plantain flour; R₉₀P₁₀ = 90% rice flour + 10% unripe plantain flour; R₈₅P₁₅ = 85% rice flour + 15% unripe plantain flour; R₈₀P₂₀ = 80% rice flour + 20% unripe plantain flour

Sensory characteristics of rice-unripe plantain *tuwo shinkafa*

Table 6 shows the mean scores for colour (5.33-7.73), taste (5.63-6.53), mouldability (6.13-6.73), aroma (4.90-6.87), texture (5.67-6.53) and overall acceptability (5.87-7.33) of rice-unripe plantain *tuwo shinkafa* based on the panellists' perceptions. The unripe plantain flour substituted *tuwo shinkafa* compared favourably with 100% rice *tuwo shinkafa* in terms of taste, mouldability, aroma, texture and overall acceptability. It was observed that while the mean scores for the colour of unripe plantain flour substituted *tuwo shinkafa* samples significantly ($p < 0.05$) decreased, their mouldability and aroma were generally enhanced. Colour is an important quality index which influences consumer's choice, preference and acceptability of food products (Akeem *et al.*, 2018). The perceived low colour quality of the unripe plantain flour substituted *tuwo shinkafa* samples by the panellists might be due to the colour of the unripe plantain which was physically observed to be yellowish compared to the whitish colour of the rice they are familiar with. The increased mouldability of the unripe plantain flour substituted *tuwo shinkafa* samples might be due to the relatively higher amylopectin to amylose ratio of unripe plantain flour compared to 100% rice

flour. Mouldability is a unique characteristic of food that enhances its swallowability at the point of consumption (Bolade and Adeyemi, 2014). Aroma is an integral part of taste and general acceptability of food. The general increase in aroma of the composite flour *tuwo shinkafa* perceived by the panellists as the level of substitution of unripe plantain flour increased might be due to the characteristic flavour of the plantain flour. This is plausible since plantains are known to have high starch content and characteristic flavour which change with acid composition during ripening (Falade and Oyeyinka, 2015). The decision to swallow or chew a food depends on its textural characteristics. No significant difference existed among the mean sensory scores obtained for the texture of *tuwo shinkafa* samples. The 10% unripe plantain substituted *tuwo shinkafa* received the highest ($p < 0.05$) overall acceptability score (7.33) among the samples. This shows that up to 20% unripe plantain could be substituted in rice flour for the production of acceptable *tuwo shinkafa* since none of the sensory qualities of the unripe plantain *tuwo shinkafa* was rated below 5 (neither like nor dislike). However, based on mean sensory ratings, 10% unripe plantain flour is recommended for substitution in rice flour for the production of *tuwo shinkafa*.

Table 6: Mean sensory scores of rice-unripe plantain *tuwo shinkafa*

Sample	Colour	Taste	Mouldability	Aroma	Texture	Overall acceptability
R ₁₀₀ P ₀	7.73±1.02 ^a	6.53±1.36 ^a	6.13±1.91 ^{ab}	4.90±1.31 ^c	6.53±1.63 ^a	6.27±1.09 ^{bc}
R ₉₅ P ₅	6.80±0.81 ^b	6.50±1.31 ^a	6.27±1.05 ^a	5.20±1.10 ^c	6.23±1.55 ^a	6.53±1.25 ^b
R ₉₀ P ₁₀	5.67±1.42 ^c	6.37±1.43 ^{ab}	6.40±1.19 ^a	5.97±1.30 ^b	5.83±1.58 ^a	7.33±1.05 ^a
R ₈₅ P ₁₅	5.33±1.24 ^c	5.67±1.37 ^b	6.47±1.07 ^a	5.97±1.38 ^b	5.73±1.39 ^a	5.97±1.04 ^{bc}
R ₈₀ P ₂₀	5.50±1.38 ^c	5.63±1.50 ^b	6.73±1.20 ^a	6.87±1.21 ^a	5.67±1.77 ^a	5.87±1.30 ^c

Values are means of thirty determinations ± SD. Means with the same superscript along the same column are not significantly ($p > 0.05$) different

R₁₀₀P₀ = 100% rice flour; R₉₅P₅ = 95% rice flour + 5% unripe plantain flour; R₉₀P₁₀ = 90% rice flour + 10% unripe plantain flour; R₈₅P₁₅ = 85% rice flour + 15% unripe plantain flour; R₈₀P₂₀ = 80% rice flour + 20% unripe plantain flour

CONCLUSION

Substitution of unripe plantain flour in rice flour enhanced the nutritional value (ash and fibre) and some functional properties (swelling capacity and water absorption capacity) of the flour blends. The microbial analysis showed increase in bacterial and fungal counts with increase in unripe plantain flour substitution and storage period. Thus, there may be need for the use of effective packaging materials to control microbial growth during long term storage of rice-unripe plantain flour blends. This study revealed that unripe plantain flour up to 20% could be substituted in rice flour for the production of highly nutritive, wholesome and acceptable *tuwo shinkafa*, though 10% substitution is recommended based on the sensory ratings.

Ethics

The study proposal was presented and then approved by the Research and Ethical committee of the Department of Home Economics and Food Science, University of Ilorin, Nigeria. The assessors are regular consumers of *tuwo shinkafa* (rice dumpling) and they gave their informed consent to participate in the sensory evaluation.

AUTHOR DISCLOSURE STATEMENT

The authors declare no conflict of interest.

REFERENCES

- Abiodun-Solanke AO, Falade KO. (2010). A review of the uses and methods of processing banana and plantain (*Musa spp.*) into storable food products. *Journal of Agricultural Research and Development*. **9**: 85–96.
- Abulude FO. (2004). Effect of processing on nutritional composition, phytate and functional properties of rice (*Oryza sativa*) flour. *Nigerian Food Journal*. **22**: 97–100.
- Adebowale AA, Sanni SA, Karim OR. and Ojoawo JA. (2010). Malting characteristics of Ofada rice: chemical and sensory qualities of malt from ofada rice grains. *International Food Research Journal*. **17**: 83–88.
- Adebowale AA, Sanni SA. and Oladapo FO. (2008). Chemical, functional and sensory properties of instant yam breadfruit flour. *Nigerian Food Journal*. **26**(1): 2–12.
- Adebowale KO. and Lawal OS (2004). Comparative study of the functional properties of Bambara groundnut (*Voandzeia subterranean*), jack bean (*Canavalia ensiformis*) and mucuna bean (*Mucuna pruriens*) flours. *Food Research International*. **37**(4): 355–365.
- Adegunwa MO, Adelekan EO, Adebowale AA, Bakare HA. and Alamu EO. (2017). Evaluation of nutritional and functional properties of plantain (*Musa paradisiaca L.*) and tigernut (*Cyperus esculentus L.*) flour blends for food formulations. *Cogent Chemistry*. **3**: 1383707.
- Adeyemi IA, Fagade SO. and Ayotade KA (1986). Some physicochemical and cooking qualities of Nigeria rice varieties. *Nigerian Food Journal*. **4**(1): 26–33.
- Agu HO and Okoli NA. (2014). Physico-chemical, sensory, and microbiological assessments of wheat-based biscuit improved with beniseed and unripe plantain. *Food Science and Nutrition*. **2**(5): 464–469.
- Akeem SA, Yerumoh O, Leigh O, Bamgbala K, Okeke G, Sokunbi F. and Olayiwola I. (2018). Physicochemical properties, colour characteristics and sensory evaluation of full cream cow-coconut milk yoghurts. *Croatian Journal of Food Science and Technology*. **10**(2): 239–244.
- Akeem SA, Mustapha BO, Ayinla RO, Ajibola O, Johnson WO. and Akintayo OA. (2023). Physical characteristics, nutritional composition and acceptability of glutenfree crackers produced from germinated pearl millet (*Pennisetum glaucum*), defatted sesame seed (*Sesamum indicum*) and defatted tigernut (*Cyperus esculentus*) composite flours. *Discover Food*. **3**(1): 22.
- Akpata MI and Akubor PI. (1999). Chemical composition and selected functional properties of sweet orange (*Citrus sinensis*) seed flour. *Plant Food for Human Nutrition*. **54**: 353–362.
- Anderson JW, Baird P, Davis RH JR, Ferreri S, Knudtson M, Koraym A, Waters V. and Williams CL. (2009). Health benefits of dietary fiber. *Nutrition Reviews*. **67**(4): 188–205.
- Anjum FM, Pasha I, Bugti MA. and Batt MS (2007). Mineral composition of different rice varieties and their milling fractions. *Pakistan Journal of Agricultural Sciences*. **44**(2): 51–58.
- AOAC. (2005). Official Methods of Analysis of the Association of Analytical Chemists International, 18th ed. (Official methods). AOAC International, Gathersburg, MD.
- Arueya GL and Akande O. (2018). Development and characterization of dumpling dough with 'optimal' dietary fibre ratio using Ofada rice (*Oryza Sativa L.*) and unripe plantain (*Musa Paradisiaca AAB*) fruit. *Integrative Food Nutrition and Metabolism*. **5**(4): 1–8.

- Aruwa CE and Akinyosoye FA. (2015). Microbiological assessment of ready-to-eat foods (Rtes) for the presence of Bacillus species. *Journal of Advances in Biology and Biotechnology*. **3**: 145-152.
- Aurore G, Berthe P. and Louis, F. (2009). Bananas, raw materials for making processed food Products: A Review. *Trends in Food Science and Technology*. **20**: 78–91.
- Ayodele OH. and Erema VG. (2010). Glycemic indices of processed unripe plantain (*Musa paradisiaca*) meals. *African Journal of Food Science*. **4(8)**: 514 - 521
- Baiyeri KP, Aba SC, Otitoju GT and Mbah OB. (2011). The effects of ripening and cooking method on mineral and proximate composition of plantain (*Musa sp AAB* cv “Agbagba”) fruit pulp. *African Journal of Biotechnology*. **10(36)**: 6979–6984.
- Bolade MK and Adeyemi IA. (2014). Quality dynamics of maize 'tuwo' (non-fermented maize-based dumpling) as influenced by steaming of maize grits at different resident time. *Journal of Food Science and Technology*. **51(11)**: 3217–3225
- Bolade MK, Usman MA, Rasheed AA, Benson EL and Salifou I. (2002). Influence of hydrothermal treatment of maize grains on the quality and acceptability of *tuwon masara* (traditional maize gel). *Food Chemistry*. **79**: 479–483.
- Chandler S. (1995) The nutritional value of bananas. In: Gowen S, editor. Bananas and Plantains. *Chapman and Hall, UK*. p. 486-480.
- Dalié DKD, Deschamps AM. and Richard-Forget F. (2010). Lactic acid bacteria–Potential for control of mould growth and mycotoxins: A review. *Food Control*. **21(4)**: 370–380.
- Falade KO and Christopher AS. (2015) Physical, functional, pasting and thermal properties of flours and starches of six Nigerian rice cultivars. *Food Hydrocolloid*. **44**: 478-490.
- Falade KO and Oyeyinka SA. (2015). Color, Chemical and Functional Properties of Plantain Cultivars and Cooking Banana Flour as Affected by Drying Method and Maturity. *Journal of Food Processing and Preservation*. **39(6)**: 816-828.
- FAO (2009). Food and Agriculture Organisation of the United Nations. Joint Meeting of the Fourth Session of the Sub-group on Bananas and the Fifth Session of the Sub-Group on Tropical Fruits held in Rome, 9–11th December 2009.
- FAO (2011). Production, commodity by country; FAO-STAT Data. Food and Agriculture Organisation of the United Nations, Rome. 2011. Available at: <http://faostat.fao.org/site/339/default.aspx>.
- Fresco L. (2005). Rice is life. *Journal of Food Composition and Analysis*. **18(4)**: 249–253.
- Giami SY and Bekebian DA. (1992). Proximate composition and functional properties of raw and processed fullfat fluted pumpkin (*Telfairia occidentalis*) seed flour. *Journal of the Science of Food and Agriculture*. **59**: 321-325.
- Inyang UE and Asuquo IE. (2016). Physico-chemical and sensory qualities of functional bread produced from wholemeal wheat and unripe plantain composite flours. *MOJ Food Processing and Technology*. **2(2)**: 48ff53.
- Juliano BO and Bechtel DB. (1985). The rice grain and its gross composition. In: Juliano BO, editor. *Rice Chemistry and Technology*. 2nd edn. The American Association of Cereal Chemists. St Paul, MI, USA. p. 17–57.
- Karim OR, Akeem SA and Arowolo TI. (2020). Effect of pretreatments and drying methods on physicochemical properties of unripe plantain flour and sensory acceptability of its cooked dough (*amala*). *Carpathian Journal of Food Science and Technology, Special Issue*. **12(5)**: 156-166.
- Kawongolo JB. (2013). Optimization of processing technology for commercial drying of bananas (Matooke). PhD dissertation. University of Kassel, Germany.
- Kolawole FL, Balogun MA, Usman FD. and Akeem SA. (2016). Effect of drying methods on the chemical and functional properties of potato (*Solanum tuberosum*) and sweet potato (*Ipomoea batatas*) varieties. *Nigerian Journal of Agriculture, Food and Environment*. **12(4)**: 151-156.
- Ogbulie JN, Ojehor SI, Isu NR. and Njoku HO. (1993). Effects of chemical and physical treatment on shelf-life of fermented African oil bean seed (*ugba*). *Nigerian Journal of Biotechnology*. **22**: 112-116.
- Ogunlakin GO, Abioye VF. and Olewepo MO. (2014). Evaluation of the quality attributes of wheat composite (wheat-cassava, wheat-plantain and wheat-rice) flours in bread making. *African Journal of Biotechnology*. **13(38)**: 3907-3911.
- Oko AO, Ubi BE, Efiue AA and Dambaba N. (2012). Comparative analysis of the chemical nutrient composition of selected local and newly introduced rice varieties grown in ebonyi state of Nigeria. *International Journal of Agriculture and Forestry*. **2(2)**: 16-23.

- Onwuka GI. (2005). Food analysis and instrumentation. Theory and Practice. Naphtali Prints, Lagos, Nigeria. p. 133–137.
- Oviasogie EF, Ehichioya O, Ogofure AG, Beshiru A, Ojo EA, Nkemakonam JO and Raphael P. (2016). Microbiological, physicochemical, and nutritional composition of plantain flour fortified with wheat flour. *International Journal of Food and Nutritional Science*. **5(3)**: 1-8.
- Sanni SA, Okeleye KA, Soyode AF. and Taiwo OC. (2005). Physicochemical properties of early and medium maturing Nigerian rice varieties. *Nigerian Food Journal*. **23**: 148-152.
- Seenaa S and Sridhar KR. (2005). Physicochemical, functional and cooking properties of under explored legumes, Canavalia of the southwest coast of India. *Food Research International*. **38**: 803–814.
- SON. (2007). Nigerian Industrial Standard for biscuits. ICS: 664.68, Standard Organization of Nigeria. p. 1-8.
- Sosulski FW, Humbert ES, Bui ES and Jones JI. (1976). Functional properties of rapeseed flours, concentrates and isolates. *Journal of Food Science*. **41**: 1349–1351.
- Tang EN, Ndindeng SA, Bigoga J, Traore K, Silue D and Futakuchi K. (2019). Mycotoxin concentrations in rice from three climatic locations in Africa as affected by grain quality, production site, and storage duration. *Food Science and Nutrition*. **00**: 1–14.
- Thomas R, Wan-Nadiah WA and Bhat R. (2013). Physicochemical properties, proximate composition, and cooking qualities of locally grown and imported rice varieties marketed in Penang, Malaysia. *International Food Research Journal*. **20(3)**: 1345-1351.
- Ukpabi UJ and Ndimele C. (1990). Evaluation of gari production in Imo State Nigeria. *Nigerian Food Journal*. **8**: 105-110.
- Umadevi M, Pushpa R, Kumar KP. and Bhowmik D. (2012). Rice-traditional medicinal plant in India. *Journal of Pharmacognosy and Phytochemistry*. **1**: 6-12.
- Verma DK and Srivastav PP. (2017). Proximate composition, mineral content and fatty acids analyses of aromatic and non-aromatic Indian rice. *Rice Science*. **24(1)**: 21–31.
- Vlachos A and Arvanitoyannis IS. (2008). A review of rice authenticity/adulteration methods and results. *Critical Review in Food Science and Nutrition*. **48**: 553–598.
- Yarkwan B and Uvir, RH. (2015). Effects of drying methods on the nutritional composition of unripe plantain flour. *Food Science and Quality Management*. **41**: 5-10.